

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/356796219>

Poor socioeconomic state, stress, stunting and fatness are negatively associated with cognitive performance of female adolescents

Article in *Sri Lanka Journal of Child Health* · December 2021

DOI: 10.4038/sljch.v50i4.9886

CITATIONS

0

READS

24

5 authors, including:



H.J.H. Madhushanthi Hemamali Madhushanthi
University of Ruhuna

12 PUBLICATIONS 19 CITATIONS

[SEE PROFILE](#)



Savithri Wimalasekera
University of Sri Jayewardenepura

78 PUBLICATIONS 87 CITATIONS

[SEE PROFILE](#)



Goonewardena Cse
University of Sri Jayewardenepura

30 PUBLICATIONS 64 CITATIONS

[SEE PROFILE](#)



Thamara Amarasekara
University of Sri Jayewardenepura

75 PUBLICATIONS 99 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Malmo OPRA study [View project](#)



Evidence Based National Framework for Undergraduate Nursing Education in Sri Lanka [View project](#)

Poor socioeconomic state, stress, stunting and fatness are negatively associated with cognitive performance of female adolescents

*H J H Madhushanthi¹, S W Wimalasekera², C S E Goonewardena³, A A T D Amarasekara⁴, Janaka Lenora⁵

Sri Lanka Journal of Child Health, 2021; 50(4): 674-680

DOI: <http://doi.org/10.4038/sljch.v50i4.9886>

Abstract

Introduction: The brain undergoes dynamic structural and maturational changes during adolescence, producing significant changes in cognition and behaviour. Cognitive performance (CP) of an individual is influenced by many factors, nutritional, psychosocial and genetic.

Objectives: To determine the influence of nutritional and psychosocial factors on CP of early female adolescents in Sri Lanka.

Method: Two hundred and eighteen female adolescents, 11-14 years of age, were administered neurocognitive test batteries comprising tests of nonverbal intelligence, subtests of Wechsler Intelligence Scale for Children and two computer based executive function tests to assess CP. Height for age Z-score (HAZ), weight for age Z-score (WAZ) and BMI for age Z-score (BAZ) were calculated to measure nutritional status. Adolescent stress questionnaire (ASQ) was used to assess stress.


Results: Fat mass percentage was inversely associated with working memory index (WMI) ($r=-0.34$; $p<0.001$) and processing speed index (PSI) ($r=-0.39$; $p<0.001$). Multivariate regression analysis revealed that the combination of socioeconomic status (SES) index, HAZ and ASQ total score significantly predicts estimated full scale intelligence quota (EFSIQ). SES index was positively correlated with verbal comprehension index (VCI) ($r=0.36$; $p<0.01$), working memory index (WMI) ($r=0.30$; $p<0.01$), EFSIQ ($r=0.31$; $p<0.01$), perceptual reasoning index (PRI) ($r=0.27$; $p<0.01$), PSI ($r=0.20$; $p<0.01$) and visuo-spatial working memory ($r=0.17$; $p<0.05$).

Conclusions: Poor socioeconomic state, stress, stunting and fatness were negatively linked with cognitive performance of female adolescents.

(Key words: Socioeconomic state, Stunting, Fatness, Cognitive performance, Female adolescents)

¹Senior Lecturer, Department of Nursing, Faculty of Allied Health Sciences, University of Ruhuna, Sri Lanka, ²Professor in Physiology, Department of Physiology, ³Professor in Community Medicine, Department of Community Medicine, Faculty of Medical Sciences, University of Sri Jayewardenepura, Sri Lanka, ⁴Senior Lecturer, Department of Nursing and Midwifery, Faculty of Allied Health Sciences, University of Sri Jayewardenepura, Sri Lanka, ⁵Professor in Physiology, Department of Physiology, Faculty of Medicine, University of Ruhuna, Sri Lanka

*Correspondence: hemamali@ahs.ruh.ac.lk

 <https://orcid.org/0000-0002-8732-9102>

(Received on 28 October 2020: Accepted after revision on 18 December 2020)

The authors declare that there are no conflicts of interest

Funding: The study was funded by the research grant from the University of Sri Jayewardenepura (ASP/06/RE/MED/2014/19).

Open Access Article published under the Creative

Commons Attribution CC-BY  License

Introduction

Ages 11 to 14 years, which is considered early adolescence, is characterized by significant physical, psychological, social and cognitive transformation¹. The prefrontal cortex of the brain, involved in high order cognitive outcomes, undergoes a different development trajectory in adolescence². As cognitive performance (CP) is influenced by nutritional, environmental and genetic factors, any insult to nutrition and environmental factors during adolescence lead to alteration of cognitive tasks governed by the frontal lobe³. In early years of life, nutrition and socio-emotional factors play a key role in cognitive development⁴. Earlier studies reported that school aged children with protein energy malnutrition had poor performance on cognitive tasks like attention, memory, executive function, visuo-spatial ability, learning and intelligence quotient^{5,6}. A multiethnic cross sectional study conducted across four South East Asian countries demonstrated that poor nonverbal intelligence scores of school aged children were associated with underweight and stunting⁷. Moreover, poor socio-economic status is associated with several cognitive deficits, language and executive function reading and school

achievements^{8,9}. In Sri Lanka, data on effect of nutritional and psychosocial factors on CP of adolescents are meagre.

Objectives

To assess nutritional and psychosocial factors affecting CP of early female adolescent school children in the Galle educational zone.

Method

This descriptive cross-sectional study was conducted on randomly selected schools in the Galle educational zone in Sri Lanka. Two hundred and eighteen 11-14 year old female adolescents residing in the Galle city were recruited. Schools and classes were randomly selected representing all school categories classified by the Ministry of Education.

Anthropometric measurements

Weight, height, waist circumference (WC) and hip circumference (HC) were measured according to standard procedure. Body mass index (BMI), fat mass percentage (FM%), waist hip ratio (WHR) and waist height ratio (WHtR) were calculated to assess nutritional status of participants. Height for age Z-score (HAZ), weight for age Z-score (WAZ) and BMI for age Z-score (BAZ) were calculated using WHO sex specific growth reference values for adolescents. Adolescents with HAZ, WAZ and BAZ reference value <-2 SD score were categorized as stunted, underweight and thinness respectively. Triceps and subscapular skinfold thickness were measured using a skinfold caliper according to standard protocol to determine body FM%. FM% was estimated using a validated fat percentage estimation equation developed for Sri Lankan children¹⁰. As WHtR does not vary with age and sex, a cutoff value could apply across all age groups, sexes and ethnic groups¹¹.

Measurement of socio-economic and psychosocial adversities

Socio-economic status (SES) is a composite index of parental education, occupation and income¹². Parental occupational status was categorized using the 9 point Hollingshead occupation scale¹³, which closely matched modern occupations. Income to need ratio was calculated by dividing total family income by official poverty threshold (published by department of Census and Statistics, Sri Lanka) for a family of that size. Subsequently, income to need ratio, parental education and Hollingshead occupation scale rank were standardized and averaged to derive family SES composite score (SES index).

The interviewer-administered psychosocial adversity scale was used to assess psychosocially adverse factors. This scale has been widely used in

previous child development studies conducted in Asia¹⁴. Adolescence stress was assessed using the interviewer-administered adolescence stress questionnaire (ASQ)^{15,16}, comprising 51 items measuring 9 dimensions of adolescence stress including stress of home life, school performance, school attendance, romantic relationships, peer pressure, teacher interaction, future uncertainty, school/leisure conflict and emerging adult responsibility. It is a self-administered questionnaire, each item rated on a 5 point Likert scale where 1= not at all stressful/ irrelevant to me, 2= a little stressful, 3= moderately stressful, 4= quite stressful and 5 = very stressful.

Measurement of cognitive performance

Three cognitive test batteries, Wechsler Intelligence tests, tests of non-verbal intelligence (TONI) and two computer based executive function tasks (inhibition and visuo-spatial working memory) were used for detailed assessment of CP. The individually administered fourth UK edition of Wechsler Intelligence Scale for Children (WISC-IV), devised by Psychological Corporation, USA, measures various cognitive skills. It is widely used to assess cognitive function of children aged 6 years to 16 years 11 months. In this study, eight subtests of WISC were used to assess four cognitive domains: verbal comprehension index (VCI) (similarities, comprehension), working memory index (WMI) (digit span, symbol search), perceptual reasoning index (PRI) (picture completion, matrix reasoning) and processing speed index (PSI) (cancellation and arithmetic). The estimated full scale intelligence quota (EFSIQ) was calculated based on the sum of the eight subtest scores. WISC profile has been validated in Sri Lanka and used in previous national studies¹⁷. TONI-3 is free of language and culture bias and an objective method of examining cognitive function which does not require verbal or motor skills. It is widely used to measure abstract reasoning of individuals aged 6 to 90 years¹⁸. It has 45 items arranged in order of difficulty per test. Executive functions were assessed using two computer based tasks, pig house paradigm and stop signal paradigm, which assess visuo-spatial memory and inhibitory control respectively.

Ethical issues: Approval for the study was obtained from the Ethics Review Committee, Faculty of Medical Sciences, University of Sri Jayewardenepura, Nugegoda, Sri Lanka (Ref. No. 80/14 dated 26 February 2015) Before starting data collection, permission was obtained from the relevant school authorities while informed written consent was obtained from parents and assent from children.

Statistical analysis: Statistical Package for Social Sciences version 20 was used to analyse data. Continuous variables not normally distributed were log transformed. Pearson correlation test was used to assess correlation of cognitive function tests scores with anthropometric and psychosocial variables. Variables with significance value <0.25 in bivariate analysis were included for regression analysis. Multivariate regression analyses were performed to predict the best combination of dependent variables on CP. In the final regression model, $p < 0.05$ was considered significant.

Results

Table 1 presents the socio-demographic characteristics of study participants. Age of study participants ranged from 11 to 14 years with a mean of 12.31 ± 1.16 years.

Table 1
Socio-demographic characteristics of participants

Variable	Frequency (%)
<i>Ethnicity</i>	
Sinhala	185 (84.9)
Muslim	33 (15.1)
<i>Religion</i>	
Buddhist	178 (81.7)
Islam	33 (15.1)
Roman Catholic	07 (03.2)
<i>Type of school</i>	
Type 2	36 (16.5)
Type C	82 (37.6)
Type 1 AB	100 (45.9)
<i>Age group (years)</i>	
11-12	104 (47.7)
12-13	75 (34.4)
13-14	39 (17.9)
<i>Family type</i>	
Nuclear	149 (68.3)
Extended	69 (31.7)
<i>Number of children in family</i>	
Single child	89 (40.8)
2-3 children	108 (49.5)
>3 children	21 (09.6)
<i>Place of residence</i>	
Urban	100 (45.9)
Semi urban	118 (54.0)
<i>Percentage of school attendance</i>	
Satisfactory	162 (74.3)
Not satisfactory	56 (25.6)
<i>Pubertal characteristics</i>	
Attained menarche	124 (56.9)
Not attained menarche	94 (43.1)

Psychological stress of adolescents

The ASQ stress score ranged from 83 to 208, mean stress score being 145.4 ± 24.3 . Whilst 158 (72.5%) belonged to the stressed group, 60 (27.5%) belonged to non-stressed group. Stress was observed for school attendance (93.1%), future uncertainty (92.7%), school performance (84.9%), emerging responsibility (85.3%) and conflict of

school/leisure time (81.7%). Most adolescents considered stress of future uncertainty and school attendance the greatest stress inducing factors.

Association of cognitive test scores and ASQ total score

Further analysis was conducted through MANOVA with group (stressed and non-stressed) as independent variables and all ten cognitive test index variables of three cognitive test batteries (WISC-IV, TONI-3 and the two computer based executive function tasks) as dependent variables. MANOVA results indicate a significant effect of stress on entire cognitive function test scores ($F(10,206) = 4.925, p < 0.001$).

Association of cognitive test scores and demographic variables

MANCOVA test was used to investigate the main effect of demographic variables on all cognitive test profile using family SES index as covariate. Effects of family type, number of siblings, place of residence, percentage of school attendance and school type on all cognitive test profiles were not statistically significant, $F(8,208) = 0.86, p > 0.05, \eta^2 = 0.03$; $F(16,414) = 1.46, p > 0.05, \eta^2 = 0.05$; $F(16,414) = 0.538, p > 0.05, \eta^2 = 0.02$; $F(16,414) = 0.538, p > 0.05, \eta^2 = 0.02$ and $F(14,414) = 1.36, p > 0.05, \eta^2 = 0.04$ respectively.

Association of cognitive test scores and socioeconomic variables

SES index was positively correlated with VCI ($r=0.36; p < 0.01$), WMI ($r=0.30; p < 0.01$), EFSIQ ($r=0.31; p < 0.01$), PRI ($r=0.27; p < 0.01$), PSI ($r=0.20; p < 0.01$) and visuo-spatial working memory ($r=0.17; p < 0.05$). Poor inhibitory control (increased stop signal delay) was associated with low SES index ($r=-0.22; p < 0.01$). Except for visuo-spatial working memory, an inverse relationship was found between PA index and scores of VCI ($r=-0.34; p < 0.01$), PSI ($r=-0.25; p < 0.01$), WMI ($r=-0.24; p < 0.01$), PRI ($r=-0.26; p < 0.01$), EFSIQ ($r=-0.34; p < 0.01$), TONI ($r=-0.13; p < 0.01$) and inhibitory control ($r=0.26; p < 0.01$).

Association of cognitive test scores and nutritional parameters

Table 2 shows the correlation between each cognitive measure and anthropometric indices. HAZ is significantly and positively associated with PSI, WMI, PRI, EFSIQ and visuo-spatial working memory. Decreased HAZ is correlated with poor inhibitory control. WAZ was not correlated with most cognitive indices except visuo-spatial working memory. No significant correlations were observed between BAZ and WHtR with any cognitive test scores.

Table 2: Correlation between cognitive test scores and anthropometric parameters

Cognitive test	HAZ	WAZ	BAZ	FM%	WC	HC	WHR	WHtR
Wechsler intelligence scale-IV								
Verbal comprehension index (VCI)	0.07 ^{ns}	0.03 ^{ns}	-0.05 ^{ns}	-0.05 ^{ns}	-0.06 ^{ns}	-0.07 ^{ns}	0.01 ^{ns}	-0.03 ^{ns}
Processing speed index (PSI)	0.28 ^{***}	0.04 ^{ns}	-0.10 ^{ns}	-0.39 ^{***}	-0.18 ^{**}	-0.15 [*]	-0.23 [*]	-0.18 ^{ns}
Working memory index (WMI)	0.18 [*]	0.04 ^{ns}	-0.12 ^{ns}	-0.34 ^{***}	-0.37 ^{***}	-0.30 ^{***}	-0.14 [*]	-0.29 ^{ns}
Perceptual reasoning index (PRI)	0.14 [*]	0.04 ^{ns}	-0.04 ^{ns}	-0.03 ^{ns}	-0.02 ^{ns}	-0.04 ^{ns}	0.04 ^{ns}	0.14 ^{ns}
Estimated full scale intelligence quota (EFSIQ)	0.22 ^{**}	0.06 ^{ns}	-0.06 ^{ns}	-0.12 ^{ns}	-0.17 [*]	-0.19 ^{**}	-0.06 ^{ns}	-0.13 ^{ns}
Test of non-verbal intelligence -3								
	-0.09 ^{ns}	0.06 ^{ns}	-0.01 ^{ns}	-0.20 ^{**}	-0.11 ^{ns}	-0.12 ^{ns}	0.05 ^{ns}	-0.02 ^{ns}
Executive function tests								
Visuo-spatial working memory	0.40 ^{***}	0.31 ^{***}	-0.02 ^{ns}	-0.21 [*]	-0.02 ^{ns}	0.02 ^{ns}	-0.05 ^{ns}	-0.16 ^{ns}
Inhibitory control	-0.22 ^{***}	-0.06 ^{ns}	-0.01 ^{ns}	0.18 [*]	0.15 [*]	0.14 [*]	0.18 [*]	-0.03 ^{ns}

ns: non-significant, *p<0.05, **p<0.01, ***p<0.001

Predictors of cognitive performance of female adolescents

School type, family type, place of residence, family size, percentage of school attendance, age group, BAZ and WHtR were not significantly associated with CP and were excluded from regression model. In bivariate analysis, anthropometric parameters (HAZ, WAZ, FM%, WC, HC, WHR) and psychosocial variables (SES index, PA index score, ASQ total score) were significantly associated with cognitive function test scores, hence qualifying for multivariable linear regression model. Data showed multi-collinearity (tolerance value <0.1) between SES index and PA index ($r=0.649$; $p<0.001$). SES

is a significant factor influencing child cognitive development. Accordingly psychosocial adversity index variable was eliminated from prediction variables and SES index included as candidate variable for multiple regression analysis.

Regression analysis was conducted to determine the extent of the variance of cognitive tests as explained by anthropometric and socioeconomic parameters. Results showed that regression model (R²) was statistically significant for the dependent variables; VCI, PRI, WMI, PSI, EFSIQ, TONI score, visuo-spatial working memory and inhibitory control (Table 3 $p<0.05$).

Table 3: Predictors of cognitive performance of female adolescents

Cognitive test	Predictor variable	Standardized Beta coefficient	p	Adjusted R ²	F	95% confidence interval
VCI	SES index	0.362	0.001*	0.131	32.47	0.25-0.52
	Constant		0.001*			
PSI	SES index	0.224	0.001*	0.163	11.51	1.78-6.14
	HAZ	0.176	0.008*			0.78-5.14
	FM%	-0.263	0.001*			-1.74- -0.60
	ASQ total score	-0.203	0.002*			-0.29 - -0.71
	Constant		0.001*			80.31-120.62
WMI	SES index	0.267	0.001*	0.200	28.19	0.16-0.43
	Waist circumference	-0.342	0.001*			-0.74- -0.35
	Constant		0.001*			101.33- 128.68
PRI	SES index	0.263	0.001*	0.101	7.06	1.22 - 3.51
	ASQ total score	-0.201	0.003*			-0.15- -0.03
	Constant		0.001*			65.61- 89.82
EFSIQ	SES index	0.256	0.001*	0.109	8.452	1.298- 3.856
	HAZ	0.211	0.001*			-1.97- -0.49
	ASQ total score	-0.181	0.006*			-0.123 - -0.021
	Constant					64.49- 71.64
TONI	FM%	-0.320	0.036	0.116	14.72	-2.73 - -0.37
	ASQ total score	-0.159	0.001*			-2.82- -0.72
	Constant		0.001*			45.36- 130.07
Visuo-spatial working memory	HAZ	0.412	0.001*	0.185	17.401	-2.224- -1.109
	SES index	0.164	0.008*			0.025-0.165
	FM%	-0.164	0.010*			0.104-0.750
	Constant		0.001*			18.68- 20.75
Inhibitory control	SES index	0.216	0.001*	0.089	11.54	-2.82- -0.72
	HAZ	0.226	0.001*			-20.82 - -5.76
	Constant		0.001*			256.95- 329.88

A combination of SES index, ASQ total score and HAZ significantly accounted for 10.9% variance of EFSIQ ($p=0.001$). SES index was found to be positively and significantly associated with most cognitive variables including VCI, PSI, WMI, PRI, EFSIQ, visuo spatial working memory and inhibitory control (Table 3). Increased HAZ was found positively associated with PSI, EFSIQ, visuo

spatial working memory and inhibitory control scores in the multivariable model. A single unit increase of FM% resulted in 0.263 unit reduction of ($\beta= -0.263$, 95% CI;-1.74- -0.60) PSI score, 0.320 unit reduction in TONI score ($\beta= -0.320$, 95% CI; - 2.73- -0.37) and 0.164 unit reduction of visuo spatial working memory ($\beta= -0.164$, 95% CI; 0.104-0.750). In the stepwise regression WAZ was

eliminated from the model due to its insignificant association after bonferroni correction.

Discussion

The present study shows that anthropometric nutritional indicators (HAZ, FM% and WC), stress, SES index and ASQ total score were significant predictors of CP amongst early female adolescents. Strengths of the present cross sectional study were the comprehensive analysis of CP using different cognitive test batteries which provide reliable findings regarding factors influencing CP. Chosen cognitive abilities are vulnerable for macronutrient deficiencies shown in previous studies^{7,19}. In line with the findings of the present study, stunting (low HAZ) is negatively associated with cognitive outcomes. Decreased availability of energy and micronutrients during the adolescent growth period resulted in decreased linear growth, reduced neurodevelopment of brain and changes in behaviour and cognition^{6,20}. Moreover, long term protein energy malnutrition leads to altered brain architecture and neuronal functions²¹. Retardation of growth is observed in the hippocampus, frontal lobe and temporal lobes which are involved in higher order cognitive tasks²². Another aspect of the role of undernutrition on deficit cognitive outcomes is explained using a social isolation theory by Grantham and Henningham²³. This theory describes that undernourished children are less explorative and apathetic with reduced activity which contribute to poor environmental stimulation which in turn leads to poor cognitive development. Impaired cognitive function in adolescence is more likely to continue into adult life and may hinder development of the full intelligence potential as an adult²⁴.

Our study found that family SES index is a robust indicator of most cognitive indices of female adolescents. This supports previous studies showing strong positive association of higher parental SES with higher scores on cognitive measures and academic tests from early childhood through adolescence^{12,24}. Evidence suggests that SES related difference in neural development begins *in utero*⁹. SES index significantly predicts verbal comprehension and executive functions (inhibitory control and visuo-spatial working memory) in this study. In a recent review, the role of underlying SES associated differences in neurocognitive abilities is attributed to multiple factors including prenatal influences, postnatal care and cognitive stimulation⁹. Inappropriate parenting practices are correlated with emotional and behavioral problems in children that even contribute to problems in later years of life²⁵.

Compared to adolescents from higher SES backgrounds, adolescents from low SES

backgrounds show more psychopathologies like internalizing (depression, schizophrenia) and externalizing behaviours²⁶. Persistent stress activates the hypothalamic-pituitary-adrenal axis and elevates stress hormones which adversely affect brain neurodevelopment and compromise cognitive and academic achievements²⁷. Continued maturation of cortico-limbic regions in adolescence makes the adolescent brain more vulnerable to stress²⁸. Some studies suggest that stress exerted structural changes in the prefrontal cortex during adolescence would cause long lasting psychopathologies compared to stress experienced during adult life²⁹. There is evidence that school based working memory training boosts the performance on mathematics, IQ and class room performance of children and adolescents³⁰. Our findings revealed that SES plays a crucial role in the neurocognitive profile of female adolescents. The study highlights the influence of low HAZ, SES, psychological stress and increased fatness indicators contribute to poor CP of female adolescents. Future research is needed to elucidate the role of fatness parameters on CP.

Conclusions

Poor socioeconomic state, stress, stunting and fatness were negatively linked with CP of female adolescents. School type, family type, place of residence, family size, percentage of school attendance, age group, BAZ and WHtR were not significantly associated with CP.

References

1. World Health Organization, (2005), 'Nutrition in adolescence: issues and challenges for the health sector: issues in adolescent health and development', Geneva. Available at: <http://www.who.int/iris/handle/10665/43342> [Accessed on 16 June 2017].
2. Sowell E. Mapping continued brain growth and gray matter density reduction in dorsal frontal cortex: inverse relationships during post- adolescent brain maturation. *Journal of Neuroscience* 2001; **21**: 8819-9929. <https://doi.org/10.1523/JNEUROSCI.21-22-08819.2001> PMID: 11698594 PMCID: PMC6762261
3. Crookston B, Penny M, Alder S, Dickerson T, Merrill R, Stanford J, *et al.* Children who recover from early stunting and children who are not stunted demonstrate similar levels of cognition. *Journal of Nutrition* 2010; **140**(11): 1996-2001.

- <https://doi.org/10.3945/jn.109.118927>
PMid: 20844188
4. Bornstein M, Putnick D. Cognitive and socioemotional caregiving in developing countries. *Child Development* 2012; **83**: 46–61.
<https://doi.org/10.1111/j.14678624.2011.01673.x>
PMid: 22277006 PMCID: PMC3270892
 5. Fink G, Rockers PC. Childhood growth, schooling, and cognitive development: further evidence from the Young Lives study. *American Journal of Clinical Nutrition* 2014; **100**(1): 182-8.
<https://doi.org/10.3945/ajcn.113.080960>
PMid: 24808488
 6. Sokolovic N, Selvam S, Srinivasan K, Thankachan P, Kurpad A, Thomas T. Catch-up growth does not associate with cognitive development in Indian school-age children. *European Journal of Clinical Nutrition* 2014; **68**(1): 14-8.
<https://doi.org/10.1038/ejcn.2013.208>
PMid: 24169458
 7. Poh B, Rojroonwasinkul N, Le Nyugen B, Budiman B, Ng L, Soonthorndhada K, et al. Relationship between anthropometric indicators and cognitive performance in Southeast Asian school-aged children. *British Journal of Nutrition* 2013; **110**(S3): S57-S64.
<https://doi.org/10.1017/S0007114513002079>
PMid: 24016767
 8. Noble K, Norman M, Farah M. Neurocognitive correlates of socioeconomic status in kindergarten children. *Developmental Science* 2005; **8**(1): 74-87.
<https://doi.org/10.1111/j.14677687.2005.00394.x>
PMid: 15647068
 9. Hackman DA, Farah MJ, Meaney MJ. Socioeconomic status and the brain: mechanistic insights from human and animal research. *Nature Reviews Neuroscience* 2010; **11**(9): 651.
<https://doi.org/10.1038/nrn2897>
PMid: 20725096 PMCID: PMC2950073
 10. Wickramasinghe V, Lamabadusuriya S, Cleghorn G, Davies P. Assessment of body composition in Sri Lankan children: Validation of a skin fold thickness equation. *Ceylon Medical Journal* 2008; **53**(3): 83-8.
<https://doi.org/10.4038/cmj.v53i3.247>
PMid: 18982800
 11. Ashwell M, Hsieh SD. Six reasons why the waist-to-height ratio is a rapid and effective global indicator for health risks of obesity and how its use could simplify the international public health message on obesity. *International Journal of Food Sciences and Nutrition* 2005; **56**(5): 303-7.
<https://doi.org/10.1080/09637480500195066>
PMid: 16236591
 12. McLoyd V. Socioeconomic disadvantage and child development. *American Psychologist* 1998; **53**(2): 185.
<https://doi.org/10.1037/0003066X.53.2.185>
PMid: 9491747
 13. Hollingshead A. Four factor index of social status. *Yale Journal of Sociology* 2011; **8**: 21–52.
 14. Bradley R, Corwyn R. Socioeconomic status and child development. *Annual Review of Psychology* 2002; **53**(1): 371-99.
<https://doi.org/10.1146/annurev.psych.53.100901.135233>
PMid: 11752490
 15. Byrne DG, Mazanov J. Adolescent stress and future smoking behaviour: A prospective investigation. *Journal of Psychosomatic Research* 2003; **54**(4): 313-21.
[https://doi.org/10.1016/S00223999\(02\)00411-7](https://doi.org/10.1016/S00223999(02)00411-7)
 16. Kathriarachchi LS, Hewage DC, Wimalasekera SW, Ranjan R, Mendis ALS. Home school environmental factors and peer pressure contributing to stress among adolescents in Colombo district in Sri Lanka – Preliminary results. *Proceedings of South Asian Association of Physiologist Conference* 2016. pp.229.
 17. Wimalasekera S. 2010. The effects of iron deficiency on cognitive function and neurophysiological function in early adolescent females. PhD Thesis. University of Sri Jayewardenepura.

18. Brown L, Sherbenou J, Johnsen S. (1990). Test of non-verbal intelligence (2nd edition). Austin, TX, USA, Pro-Ed.
19. Eilander A, Muthayya S, van der Knaap H, Srinivasan K, Thomas T, Kok F, et al. Undernutrition, fatty acid and micronutrient status in relation to cognitive performance in Indian school children: a cross-sectional study. *British Journal of Nutrition* 2010; **103**(7): 1056-64.
<https://doi.org/10.1017/S000711450999273X>
PMid: 20003612
20. Kar B, Rao S, Chandramouli B. Cognitive development in children with chronic protein energy malnutrition. *Behavioral and Brain Functions* 2008; **4**(1): 31.
<https://doi.org/10.1186/1744-9081-4-31>
PMid: 18652660 PMCID: PMC2519065
21. Bryan J, Osendarp S, Hughes D, Calvaresi E, Baghurst K, van Klinken J. Nutrients for cognitive development in school-aged children. *Nutrition Reviews* 2004; **62**(8): 295-306.
<https://doi.org/10.1111/j.17534887.2004.tb00055.x>
PMid: 15478684
22. Benton D, Winichagoon P, Ng T, Tee E, Isabelle M. Symposium on nutrition and cognition: towards research and application for different life stages. *Asia Pacific Journal of Clinical Nutrition* 2012; **12**: 104-24.
23. Grantham-McGregor S, Baker-Henningham H. Review of the evidence linking protein and energy to mental development. *Public Health Nutrition* 2005; **8**(7a): 1191-201.
<https://doi.org/10.1079/PHN2005805>
PMid: 16277829
24. Hanscombe K, Trzaskowski M, Haworth C, Davis O, Dale P, Plomin R. Socioeconomic status and children's intelligence: In a UK-representative sample SES moderates the environmental, not genetic, effect on IQ. *PLoS One* 2012; **7**(2): e30320.
<https://doi.org/10.1371/journal.pone.0030320>
PMid: 22312423 PMCID: PMC3270016
25. sWadsworth M, Raviv T, Compas B, Connor-Smith J. Parent and adolescent responses to poverty related stress: Tests of mediated and moderated coping models. *Journal of Child and Family Studies* 2005; **14**(2): 283-98.
<https://doi.org/10.1007/s10826-005-5056-2>
26. Qin S, Hermans E, van Marle H, Luo J, Fernández G. Acute psychological stress reduces working memory-related activity in the dorsolateral prefrontal cortex. *Biological Psychiatry* 2009; **66**(1), 25-32.
<https://doi.org/10.1016/j.biopsych.2009.03.006>
PMid: 19403118
27. Arnsten A, Li B. Neurobiology of executive functions: catecholamine influences on prefrontal cortical functions. *Biological Psychiatry* 2005; **57**(11): 1377-84.
<https://doi.org/10.1016/j.biopsych.2004.08.019>
PMid: 15950011
28. Romeo R. The impact of stress on the structure of the adolescent brain: implications for adolescent mental health. *Brain Research* 2017; **1654**: 185-91.
<https://doi.org/10.1016/j.brainres.2016.03.021>
PMid: 27021951
29. Walker S, Wachs T, Gardner J, Lozoff B, Wasserman G, Pollitt E, et al. Child development: risk factors for adverse outcomes in developing countries. *Lancet* 2007; **369**(9556): 145-57.
[https://doi.org/10.1016/S01406736\(07\)60076-2](https://doi.org/10.1016/S01406736(07)60076-2)
30. St Clair-Thompson H, Stevens R, Hunt A, Bolder E. Educational Psychology: An International Journal of Experimental Educational Psychology. *Educational Psychology* 2010; **30**(2): 203-19.
<https://doi.org/10.1080/01443410903509259>