

TABLE OF CONTENTS

	Page
Acknowledgements	i
Abstract	ii
Table of contents	v
List of Figures	xiii
List of Tables	xvii
Chapter 1	
Introduction	1
Chapter 2	
Outline of the review of the literature	3
Chapter 3	
Factors affecting the fluid exchange across the capillary wall - - Starling forces	4
3.1 Sequence of liquid accumulation during formation of pulmonary oedema	4
3.2 Structure of the alveolar-capillary membrane	5
3.2.1 Capillary endothelial cell layer	5
3.2.2 The interstitial space	6
3.2.3 The lining of the alveolar wall	6
3.3 Bronchial Circulation	7
3.4 Starling forces	8
3.5 Factors affecting the accumulation of interstitial fluid in the lungs	11

3.6	Methods of increasing the extravascular fluid content in the lungs of experimental animals	13
Chapter 4		
	Review on pulmonary sensory receptors	15
4.1	Slowly adapting receptors (SAR)	15
4.2	Rapidly adapting receptors (RAR)	20
4.3	C fibre afferent receptors	24
Chapter 5		
	Experimental models of chronic heart failure	30
5.1	Models of chronic mitral regurgitation	30
5.2	Models involving aortic valve or aorta	32
5.3	Models involving pulmonary valve or pulmonary artery	33
5.4	High output cardiac failure caused by aorto-caval fistula	33
5.5	Models of heart failure caused by rapid pacing of the heart	34
5.6	Doxorubicin (adriamycin) induced heart failure	34
5.7	Heart failure secondary to myocardial infarction	35
Chapter 6		
	Methods of assessing lung water	36
6.1	Direct methods	36
6.2	Indirect methods	37
Chapter 7		
	Statement of the problem	39

Chapter 8

Methods	41
8.1 Surgical induction of mitral regurgitation	41
8.2 General surgical preparation	46
8.3 Electrophysiological studies	47
8.4 Measurement of conduction velocity of the axons	48
8.5 Identification of nerve fibres arising from RAR, SAR and C fibre receptors in the lung and the airways	49
8.6 Location of the receptors	50
8.7 <i>In vivo</i> assessment of mitral regurgitation, left ventricular hypertrophy and left ventricular function: Echocardiography	51
8.8 Statistical analysis	52

Chapter 9

General results	53
-----------------	----

Chapter 10

Evidence for mitral valve damage	59
10.1 Visual inspection of the mitral valve apparatus	59
10.2 Post mortem evidence for left ventricular hypertrophy: increased left ventricular weight	59
10.3 Evidence for pulmonary venous congestion: elevated left atrial pressure	59
10.4 <i>In vivo</i> assessment of mitral regurgitation, left ventricular hypertrophy and left ventricular function: echocardiography	63

Chapter 11

Effect of acute elevation of left atrial pressure to different degrees on the activity of pulmonary vagal afferents	67
11.1 Acute elevation of left atrial pressure	67
11.2 Experimental Protocols	69
11.2.1 Protocol 1a: Effect of acute graded elevation of left atrial pressure on the activity of vagal afferents	69
11.2.2 Protocol 1b: Effect of acute elevation of left atrial pressure above 25 mmHg to induce pulmonary oedema	70
11.3 Results	71
11.3.1 Protocol 1a: Effect of acute graded elevation of left atrial pressure on the activity of vagal afferents	71
11.3.1.1 Intact control rabbits	71
11.3.1.1.1 RAR	71
11.3.1.1.2 SAR	74
11.3.1.2 Sham operated rabbits	77
11.3.1.2.1 RAR	77
11.3.1.3 Rabbits with chronic pulmonary venous congestion for 6 weeks	79
11.3.1.3.1 RAR	79
11.3.1.4 Rabbits with chronic pulmonary venous congestion for 12 weeks	81
11.3.1.4.1 RAR	81

11.3.1.4.2	SAR	88
11.3.1.4.3	Bronchial C fibre afferents	91
11.3.1.4.4	Pulmonary C fibre afferents	94
11.3.2	Protocol 1b: Effect acute elevation of left atrial pressure above 25 mmHg on the activity of vagal afferents	97
11.3.2.1	Rabbits with chronic pulmonary venous congestion for 12 weeks	97
11.3.2.1.1	RAR	97
11.3.2.1.2	C fibre afferents	100

Chapter 12

Protocol 2:	Effect of obstruction of pulmonary lymphatic drainage on the activity of pulmonary vagal afferents	103
12.1	Experimental protocols	
12.1.1	Effect of obstruction of lymph drainage from the lung on C fibre afferent and RAR activity	105
12.1.2	Effect of combining obstruction of lymph drainage from the lung and plasmapheresis on C fibre afferent and RAR activity	105
12.2	Results	105
12.2.1	Effect of pulmonary lymphatic obstruction and plasmapheresis on pulmonary vagal afferents	106
12.2.1.1	Intact control rabbits	106
12.2.1.1.1	RAR	106

12.2.1.1.2	Pulmonary C fibre afferents	110
12.2.1.1.3	Bronchial C fibre afferents	115
12.2.1.2	Rabbits with chronic pulmonary venous congestion for 12 weeks	118
12.2.1.2.1	RAR	118
Chapter 13		
Protocol 3: measurement of the water content in the trachea, bronchi, lungs and the left ventricle		120
13.1	Results	122
Chapter 14		
Discussion		126
14.1	Selection of the species	127
14.2	Rabbits with chronic mitral regurgitation	128
14.3	Pulmonary vagal afferents in pulmonary venous congestion and oedema	130
14.3.1	RAR	130
14.3.2	SAR	133
14.3.3	Bronchial C fibre afferents	134
14.3.4	Pulmonary C fibre afferents	135
14.3.5	Haemodynamic response of rabbits during acute elevation of LAP	137
14.4	Pulmonary vagal afferents and obstruction of pulmonary lymphatic drainage	137

14.4.1	RAR	137
14.4.2	Pulmonary C fibre afferents	139
14.5	Alteration of Starling forces and extravascular water content in the lungs and the airways	139
14.5.1	Lung tissues	139
14.5.2	Major airways	141
 Chapter 15		
Conclusions		143
15.1	RAR	143
15.2	SAR	143
15.3	Pulmonary C fibre afferents	143
15.3	Bronchial C fibre afferents	144
15.4	Extravascular water content	144
 Chapter 16		
Clinical implications		145
References		147
Appendix 1:	Chemicals, materials, instruments and equipment	
Appendix 2:	Calculation of extravascular water content in the tissues	
Appendix 3:	Publications	
3.1	Gunawardena S., Bravo E., Kappagoda C. T. (1998). Effect of chronic mitral valve damage on activity of pulmonary rapidly adapting receptors in the rabbit. <i>Journal of Physiology</i> 511.1, 79-88.	
3.2	Gunawardena S., Bravo E., Kappagoda C. T. (1999). Rapidly adapting receptors in a rabbit model of mitral regurgitation. <i>Journal of Physiology</i> 521.3, 739-748.	
3.3	Symons J. D., Gunawardena S., Kappagoda C. T., Dhond M. R. (2001). Volume	

overload left ventricular hypertrophy: effects on coronary microvascular reactivity in rabbits. *Experimental Physiology* 86.6, 725-732.

- 3.4 Gunawardena S., Ravi K., Longhurst J. C., Kaufman M. P., Ma Amy, Bravo Milo, Kappagoda C. T. (2002). Response of C fibre afferents of the rabbit airways and lungs to changes in extra-vascular fluid volume. *Respiratory Physiology and Neurobiology* 132, 239-251

List of Figures

No.	Title	Page
Fig. 1	Photograph of the micrscissors used to damage the mitral valve leaflets.	43
Fig. 2	Schematic diagram showing the procedure of mitral valve damage.	44
Fig. 3	Example of a slowly adapting receptor (SAR).	54
Fig. 4	Example of a rapidly adapting receptor (RAR).	54
Fig. 5	Example of a pulmonary C fibre afferent receptor.	55
Fig. 6	Example of a bronchial C fibre afferent.	55
Fig. 7	Pictures of mitral valve apparatus of rabbits. Panel A shows a normal valve leaflets. Panels B, C and D show examples of perforated valve leaflets.	60
Fig. 8	Echocardiograms of rabbits with chronic pulmonary venous congestion showing 1+ (top) and 4+ (bottom) mitral regurgitation.	64
Fig. 9	Activity of RAR in response to acute graded elevation of LAP in control rabbits.	72
Fig. 10	Activity of an RAR in a control rabbit during acute graded elevation of left atrial pressure.	73
Fig. 11	SAR activity in control rabbits in response to acute graded elevation of LAP.	75
Fig. 12	Activity of an SAR in a control rabbit during acute graded elevation of LAP.	76
Fig. 13	Response of RARs to acute graded elevation of LAP in sham operated rabbits.	78
Fig. 14	Activity of RARs in rabbits with chronic pulmonary venous	

congestion for 6 weeks.	80
Fig. 15 Activity of RAR in rabbits with chronic pulmonary venous congestion for 12 weeks.	82
Fig. 16 Response of an RAR to acute graded elevation of LAP in a rabbit with chronic pulmonary venous congestion for 12 weeks.	83
Fig. 17 Activity of RARs in rabbits with chronic pulmonary venous congestion for 12 weeks (a) before and (b) after hypertonic albumin infusion.	85
Fig. 18 RAR activity in a rabbit with chronic pulmonary venous congestion for 12 weeks, showing the response to acute graded elevation of LAP (A) before and (B) after an intravenous infusion of hypertonic albumin solution.	86
Fig. 19 SAR activity in rabbits with chronic pulmonary venous congestion for 12 weeks.	89
Fig. 20 Activity of an SAR in a rabbit with chronic pulmonary venous congestion, during acute graded elevation of LAP.	90
Fig. 21 Activity of bronchial C fibre afferents in rabbits with chronic pulmonary venous congestion for 12 weeks.	92
Fig. 22 Effect of acute graded elevation of LAP on the activity of a bronchial C fibre afferent in a rabbit with chronic pulmonary venous congestion.	93
Fig. 23 Activity of pulmonary C fibre afferents in rabbits with chronic pulmonary venous congestion for 12 weeks.	95
Fig. 24 Response of pulmonary C fibre afferent to acute graded elevation of LAP in a rabbit with chronic pulmonary venous congestion for 12 weeks.	96

Fig. 25	Response of RARs in rabbits with chronic pulmonary venous congestion for 12 weeks during acute elevation of LAP by 25 mmHg.	98
Fig. 26	Response of an RAR to acute elevation of LAP above 25 mmHg in a rabbit with chronic pulmonary venous congestion for 12 weeks.	99
Fig. 27	Activity of C fibre afferents in rabbits with chronic pulmonary venous congestion for 12 weeks during acute elevation of LAP by 25 mmHg.	101
Fig. 28	Response of a bronchial C fibre afferent to acute elevation of LAP above 25 mmHg in a rabbit with chronic pulmonary venous congestion for 12 weeks.	102
Fig. 29	Diagram of the experimental preparation which was used to obstruct the pulmonary lymphatic drainage	104
Fig. 30	Activity of RARs in response to lymphatic obstruction and plasmapheresis in intact control rabbits.	108
Fig. 31	Example of a RAR activity in response to pulmonary lymphatic obstruction before and after plasmapheresis in a control rabbit.	109
Fig. 32	Activity of pulmonary C fibre afferents in control rabbits when pulmonary lymphatic drainage was obstructed.	111
Fig. 33	Activity of pulmonary C fibre afferents in response to pulmonary lymphatic obstruction, before (a) and after (b) plasmapheresis.	113
Fig. 34	Example of a pulmonary C fibre activity in response to pulmonary lymphatic obstruction before and after plasmapheresis in a control rabbit.	114
Fig. 35	Activity of bronchial C fibre afferents in response to PLO.	116
Fig. 36	Activity of bronchial C fibre afferents in response to PLO before and after plasmapheresis.	117

Fig. 37 RAR activity in rabbits with chronic pulmonary venous congestion for
12 weeks in response to pulmonary lymphatic obstruction.

119

List of Tables

No.	Title	Page
Table 1	Nerve conduction velocities (m s^{-1}).	56
Table 2.	Stimulus parameters used in different types of vagal pulmonary receptors when measuring the nerve conduction velocity of their axons.	57
Table 3.	Heart rate, mean arterial blood pressure, mean left atrial pressure and peak intratracheal pressure in different groups of rabbits at the beginning of experiments.	58
Table 4	Left ventricular weights and body weights	61
Table 5	Left atrial pressures	62
Table 6	Echocardiographic indices of left ventricular hypertrophy	65
Table 7	Plasma albumin, total protein, plasma osmolality, plasma sodium and haematocrit values before and after the infusion of hypertonic albumin solution.	87
Table 8	Total water content in major airways, lungs and the left ventricle (as a % of wet weight).	123
Table 9	Extravascular water content in major airways, lungs and the left ventricle (as % of wet weight).	124
Table 10	Cardiorespiratory parameters in rabbits whose tissues were harvested for analysis of water content.	125