

Contents

| | |
|---|------|
| Declaration of Authorship | i |
| Acknowledgements | ii |
| List of Figures | viii |
| List of Tables | xii |
| Abbreviations | xiii |
| Notations | xiv |
| Abstract | xv |
| | |
| 1 Introduction | 1 |
| 1.1 What is pattern formation? | 1 |
| 1.2 Overview | 2 |
| 1.3 Physiology of corals | 3 |
| 1.3.0.1 How coral polyps feed | 4 |
| 1.3.1 Reproduction process of polyps | 5 |
| 1.3.2 Coral Identification | 5 |
| 1.3.2.1 Coral morphology | 6 |
| 1.3.3 Stony corals | 8 |
| 1.3.3.1 LPS corals | 9 |
| 1.3.3.2 SPS corals | 9 |
| 1.3.4 Coral patterns and growth factors | 10 |
| | |
| 2 Modelling pattern formation process | 13 |
| 2.1 Introduction to Reaction, Reaction-Diffusion, Reaction-Diffusion-Advection Systems | 13 |
| 2.2 Pattern Formation and Reaction-Diffusion Systems | 15 |
| 2.2.1 Turing instability | 16 |
| 2.2.2 Some pattern formation Reaction-Diffusion systems | 20 |

| | | |
|----------|---|-----------|
| 2.2.2.1 | Alan Turing's spot-formation system | 20 |
| | | 20 |
| | | 20 |
| 2.2.2.2 | Gray-Scott model | 21 |
| 2.2.2.3 | Hans Meinhardt's spot-formation system | 22 |
| 2.2.2.4 | Hans Meinhardt's strip-formation system | 23 |
| 2.2.3 | Branching patterns by RDS | 23 |
| 2.2.3.1 | Spatial distribution patterns of bacterial | 23 |
| 2.2.3.2 | Some other branching structures by RDS | 26 |
| 2.3 | Pattern Formation by L-Systems and Fractals | 26 |
| 2.3.1 | L-Systems | 26 |
| 2.3.2 | Fractals | 29 |
| 2.4 | Branching structures by diffusion limited aggregation(DLA) method | 30 |
| 3 | Dynamics of ODEs | 32 |
| 3.0.1 | Autonomous system | 32 |
| 3.0.2 | Non-autonomous system | 33 |
| 3.1 | Stability of the solutions of ODE systems | 33 |
| 3.1.1 | Poincarè stability | 34 |
| 3.1.2 | Liapunov stability | 34 |
| 3.1.3 | Stability and linear approximation in two dimensions | 35 |
| 3.1.4 | Local stability via linear approximations | 36 |
| 3.1.5 | global stability | 37 |
| 3.1.5.1 | Liapunove functions and global stability | 37 |
| 3.2 | Limit Cycles | 38 |
| 3.3 | Bifurcation analysis | 38 |
| 3.3.1 | Local bifurcation | 39 |
| 3.3.2 | Hopf Bifurcation Analysis | 39 |
| 3.3.2.1 | Supercritical Hopf bifurcation | 40 |
| 3.3.2.2 | Subcritical Hopf bifurcation | 41 |
| 3.3.3 | Saddle-node bifurcation | 41 |
| 3.3.4 | Global bifurcations | 42 |
| 4 | Formulation of the models | 43 |
| 4.1 | Introduction to Modelling | 43 |
| 4.1.1 | Mathematical modelling | 43 |
| 4.2 | Derivation of a mathematical model for coral growth | 44 |
| 4.2.1 | Chemical reaction process of coral formation | 44 |
| 4.2.1.1 | Coral's nutrients and the role of the $C_6H_{12}O_6$ of the above chemical reaction process | 45 |
| 4.3 | Formulation of some models | 46 |
| 4.3.1 | Assumptions and limitations | 46 |
| 4.3.2 | A reaction type mathematical model | 47 |

| | | |
|----------------------------|---|-----------|
| 4.3.2.1 | Nondimensionalization | 48 |
| 4.3.3 | Another form of reaction type model | 49 |
| 4.3.3.1 | Nondimensionalization | 50 |
| 4.3.4 | A Reaction-Diffusion Type mathematical model | 50 |
| 4.3.5 | Simplification to a two-dimensional model | 51 |
| 4.3.6 | Nondimensionalization | 52 |
| 4.3.7 | A Reaction-Diffusion-Advection Type mathematical model | 52 |
| 4.3.8 | Simplification to a two-dimensional model | 53 |
| 4.3.9 | Nondimensionalization | 55 |
| 4.3.10 | Modifying the model incorporate the aggregation of solid material on coral surface | 55 |
| 5 | Stability of equilibria | 57 |
| 5.1 | Steady states | 57 |
| 5.1.1 | Case I ($\alpha > 2\lambda$): | 58 |
| 5.1.2 | The local behavior of the equilibrium points via linearization | 60 |
| 5.1.2.1 | The linear stability of the steady state S_1 | 60 |
| 5.1.2.2 | The linear stability of the steady state S_2 | 60 |
| 5.1.2.3 | The linear stability of the steady state S_3 | 62 |
| 5.1.3 | The structure of the trajectories in a region containing all the steady states(Global behavior) | 62 |
| 5.1.3.1 | Heteroclinic connections and phase diagram in parameter regions <i>I</i> and <i>II</i> | 66 |
| 5.1.3.2 | Heteroclinic connections and phase diagrams in parameter region <i>III</i> and <i>IV</i> | 69 |
| 5.1.3.3 | Seperatrixes | 70 |
| 5.1.4 | Existence of oscillatory solutions | 72 |
| 5.1.4.1 | Stable and unstable trajectories and phase diagrams about a neighborhood of Hopf bifurcation line | 73 |
| At parameter point P_0 : | | 73 |
| At parameter point P_1 : | | 74 |
| At parameter point P_2 : | | 75 |
| 5.1.5 | Existence of limit cycles | 78 |
| 5.1.5.1 | Limit cycles in the Region III | 78 |
| 5.1.5.2 | Growth forms corresponding to initial state lie inside and out sides of the limit cycle | 78 |
| 5.1.6 | Case II: $\alpha = 2\lambda$ | 80 |
| 5.1.7 | Case III: $\alpha < 2\lambda$ | 81 |
| 6 | Pattern formation by RDS | 84 |
| 6.1 | Turing instability region | 85 |
| 6.2 | Range of unstable wave numbers | 89 |
| 6.3 | Development of the branching structure | 99 |
| 6.3.1 | Aggregation of Calcium Carbonate on of corals | 99 |

| | |
|---|------------|
| 7 Numerical results | 103 |
| 7.1 RDS with time independent parameters | 103 |
| 7.1.1 One dimensional case | 103 |
| 7.1.2 Two dimensional case | 106 |
| 7.2 RDS with time dependent parameters | 112 |
| 7.2.1 Pattern for linearly time dependent α ($\alpha \propto t$) | 112 |
| 7.2.2 Patterns for quadratically time dependent α ($\alpha \propto t^2$) | 113 |
| 7.2.3 Patterns for $\alpha \propto t^4$ | 115 |
| 8 Existence of the stationary solutions | 116 |
| 8.1 Priori estimates | 117 |
| 8.2 Nonexistence of non constant solutions | 119 |
| 8.3 Existence of non constant solutions | 124 |
| 8.3.1 The case $\alpha = 2\lambda$ | 126 |
| If $\mu_m = \mathbf{0}$: | 127 |
| If $\mu_m > \mathbf{0}$: | 127 |
| 8.3.2 The case $\alpha > 2\lambda$ | 128 |
| 8.3.2.1 The case $\mathbf{w}_* \equiv (u_2, v_2)$ | 128 |
| If $\mu_m = \mathbf{0}$: | 128 |
| If $\mu_m > \mathbf{0}$: | 128 |
| 8.3.2.2 The case $\mathbf{w}_* \equiv (u_3, v_3)$ | 129 |
| If $\mu_m = \mathbf{0}$: | 129 |
| If $\mu_m > \mathbf{0}$: | 130 |
| 9 Discussion | 132 |
| 9.1 Future work | 135 |
| 9.1.1 Analytical Investigations | 135 |
| 9.1.2 Numerical work | 136 |
| 9.1.3 Constructing new models | 136 |
| 9.1.4 JAVA programs for simulations and visualization | 136 |
| A Important results from Functional Analysis | 137 |
| A.1 Maximum principle and some inequalities | 137 |
| Maximum principle ([1]) | 137 |
| Harnack inequality ([1]) | 138 |
| A.2 Topological Degree Theories | 138 |
| B Finite difference schemes for RDS | 140 |
| C C Codes | 145 |
| D Matlab Codes | 160 |
| D.1 MATLAB code for solving one dimensional RDS and plotting the results . | 160 |

| | |
|--|------------|
| D.2 MATLAB code for density plot | 162 |
| D.3 MATLAB code for 3D-visualization | 164 |
| | |
| Bibliography | 164 |

List of Figures

| | | |
|-----|--|----|
| 1.1 | A section of a coral polyp: Reproduced from [2] | 4 |
| 1.2 | Different types of corals | 7 |
| 1.3 | Some branching corals. Reproduced from [3] | 8 |
| 1.4 | LPS corals: Reproduced from [4] | 9 |
| 1.5 | SPS corals:reproduced from [4] | 10 |
| 1.6 | Three specimens of the coral <i>pocillopora damicornis</i> from sites increasingly sheltered from water movement. Specimen (a) is most exposed to flow; specimen (c) is most protected from flow. Reproduced from [5] | 12 |
| 2.1 | Computer simulated bacteria spread patterns: Reproduced from [6] | 25 |
| 2.2 | Reproduced from [7] | 28 |
| 2.3 | Tree like structure: Reproduced from [8] | 29 |
| 2.4 | Reproduced from [9] | 30 |
| 2.5 | Computer simulated branching coral like structures: Reproduced from [10] | 31 |
| 3.1 | General transition between types of the stable point-Reproduced as in [11] | 36 |
| 3.2 | Possible types of supercritical Hopf bifurcations in two dimension | 41 |
| 3.3 | Possible types of subcritical Hopf bifurcations in two dimension | 41 |
| 4.1 | Information flow in model building [12] | 44 |
| 4.2 | Sketch of the reaction process of the model system | 47 |
| 4.3 | Sketch of the reaction process of the model system | 49 |
| 4.4 | | 54 |
| 5.1 | Sketch of the variation of u_s and v_s with respect to α : (a) u_s ; (b) v_s | 58 |
| 5.2 | Nullclines and steady states of (5.1) for $\alpha = 10$, $\lambda = 4$ (the case $\alpha > 2\lambda$) | 59 |
| 5.3 | Plots of u_s and v_s for different values of λ (from left curve to right curves, values of λ are 1, 3, 5, 7 and 9 respectively). (a) Solid and dotted curve represent u_{s2} and u_{s3} respectively; horizontal solid line is u_{s1} , (b) Solid and dotted curve represent v_{s2} and v_{s3} respectively; Horizontal solid line (along α axis) is v_{s1} | 59 |
| 5.4 | Different regions of the (α, λ) space corresponding to equilibrium point S_2 . | 61 |

| | | |
|------|--|----|
| 5.5 | Behavior of the linearized system about S_2 at parameter points which lie in different parameter regions. (a) At region I: $\alpha = 25$, $\lambda = 12$, in this region S_2 is an unstable node. (b) At region II: $\alpha = 24$, $\lambda = 10$, in this region S_2 is an unstable spiral. (c) At region III: $\alpha = 20$, $\lambda = 6$, In this region S_2 is an stable spiral. (d) At region IV: $\alpha = 12.5$, $\lambda = 2$. In this region S_2 is stable node. | 63 |
| 5.6 | Behaviors of the linearized system about S_2 at two parameter points (a) on Hopf bifurcation curve for $\lambda = 3.5$; (b) on the line $\alpha = 2\lambda$ for $\lambda = 1.5$ | 64 |
| 5.7 | (a)The region \mathcal{R} ; (b) A confined set containing S_1 , S_2 and S_3 | 65 |
| 5.8 | Stable and unstable trajectories starting at S_3 : (a) in parameter region I for $\alpha = 25$, $\lambda = 12$, (b) in parameter region II for $\alpha = 24$, $\lambda = 10$ | 67 |
| 5.9 | A phase plane diagram (a) at region I for $\lambda = 12$, $\alpha = 25$; (b) at region II for $\lambda = 10$, $\alpha = 24$ | 67 |
| 5.10 | (a): some initial states in parameter region I which lie on stable and unstable trajectories at S_3 and some other initial states. Growth forms corresponding to initial states (b) A_1 , A_2 , A_3 , A_4 and (c) B_1 , B_2 , B_3 and B_4 | 68 |
| 5.11 | Growth forms with an initial state near S_2 at region I and II . $C1$ and $C2$ denote the growth forms in region I and II respectively | 69 |
| 5.12 | Stable and unstable trajectories starting at S_3 in region: (a) III for $\alpha = 20$, $\lambda = 6$; (b) IV for $\alpha = 12.5$, $\lambda = 2.0$; (c) IV for $\alpha = 2.4$, $\lambda = 0.5$ | 70 |
| 5.13 | Phase plane diagram for: (a) region III for $\lambda = 6$, $\alpha = 20$;(b)region IV for $\lambda = 2$, $\alpha = 12.5$. Domain of attraction (Shaded areas) of S_1 in regions: (c) III for $\lambda = 6$, $\alpha = 20$ and (d) IV for $\lambda = 2$, $\alpha = 12.5$ | 71 |
| 5.14 | (a) Some initial states in the parameter region III which are lying on stable and unstable trajectories at S_3 and some other initial states. Growth forms corresponding to initial states: (b) A_1 , A_2 , A_3 , A_4 and (b) B_1 , B_2 , B_3 and B_4 | 72 |
| 5.15 | Three points in a neighborhood of Hopf-bifurcation line | 73 |
| 5.16 | (a) Stable and unstable trajectories starting at S_3 and (b) phase plane diagram corresponding to parameter point P_0 for $\lambda_0 = 3.5$ | 74 |
| 5.17 | Growth forms at P_0 when the initial state lies: (a) above the seperatrix (b) on the seperatrix (c) below the seperatrix. | 74 |
| 5.18 | (a) stable and unstable trajectories starting at S_3 and (b) phase plane diagram; corresponding to parameter point P_1 , for $\lambda_0 = 3.5$, $\delta = 0.009$ | 75 |
| 5.19 | Growth forms at P_1 when initial state lies: (a) on the stable trajectory of S_3 , which starts at S_2 (b) near the S_2 which doesn't lie on stable trajectory of S_3 (c) sufficiently away from S_2 | 75 |
| 5.20 | Stable and unstable trajectories of S_3 corresponding to parameter point P_2 , for $\lambda_0 = 3.5$, $\delta = 0.009$ | 76 |
| 5.21 | (a) Phase plane diagram (b) Phase plane diagram in a neighborhood of S_2 ; corresponding to parameter point P_2 for $\lambda_0 = 3.5$, $\delta = 0.009$ | 76 |

| | |
|---|-----|
| 5.22 Growth forms at P_2 when the initial state lies (a) on a point of the stable trajectory of S_3 which starts at S_2 (b) at a point inside the limit cycle (c) at a point near the S_2 (Outside the limit cycle) which doesn't lie on stable trajectory of S_3 (d) at a point sufficiently away from S_2 | 77 |
| 5.23 Stable and unstable trajectories at S_3 at parameter point $((p_2)\alpha_1 + \delta, \lambda)$ when $\lambda = 3.5$ for: (a) $\delta = 0.001$ (b) $\delta = 0.01$ (c) $\delta = 0.05$ and (d) $\delta = 0.095$ | 79 |
| 5.24 Homoclinic connection at S_3 for $\lambda = 3.5$, $\delta = \delta_c(3.5) = 0.098$ | 79 |
| 5.25 Three initial points inside and out side of the limit cycle. | 80 |
| 5.26 Growth forms corresponding A , B and C | 80 |
| 5.27 Nullclines for $\alpha = 10$, $\lambda = 5$ (the case $\alpha = 2\lambda$). | 80 |
| 5.28 Phase plane diagram for the case $\alpha = 2\lambda$: (a) when $\lambda = 1.5$ (b) when $\lambda = 2$ (c) when $\lambda = 3$ | 81 |
| 5.29 (a) nullclines and (b) phase plane diagram for $\alpha = 8$, $\lambda = 5$ (a parameter point in region $\alpha < 2\lambda$) | 82 |
| 6.1 Curves to determine Turing space for $d = 0.001$ | 86 |
| 6.2 The curve $\alpha = (TC2)\alpha_1$ and $\alpha = (TC4)\alpha_1$ curves for different values of d . . | 87 |
| 6.3 λ_1 and λ_2 curves with respect to d | 88 |
| 6.4 Sketches of Turing space. In each figure dashed curve is $\alpha = (TC2)\alpha_1$, solid curve is $\alpha = (TC4)\alpha_1$ and the straight line is $\alpha = 2\lambda$. Shaded area is the Turing space. | 88 |
| 6.5 (λ, α) parameter space(shaded area) for different vales of d . In each Figure, upper curve(dashed curve): $\alpha = (TC_4)\alpha_1$, middle curve(dotted curve): $\alpha = (TC_2)\alpha_1$, lower line(solid line): $\alpha = 2\lambda$ | 89 |
| 6.6 The plot of d_c verses α for different values of λ | 90 |
| 6.7 The plot of $\max(d_c)$. The dotted line is $d = (3 - 2\sqrt{2})$ | 91 |
| 6.8 The plot of $h(\mu)$ for $d = d_c = 0.648532$, $d = 0.6 < d_c$, $d = 0.4 < d_c$ and $d = 0.75 > d_c$ | 92 |
| 6.9 The plot of σ_1 agains μ for some different values of σ when $\lambda = 3$ | 93 |
| 6.10 The plot of $\max\sigma_1$; from left to right curves are for $\lambda = 2$, $\lambda = 6$, $\lambda = 9$ respectively | 94 |
| 6.11 The plot of $n_{\max\sigma}$; from left to right curves are for $\lambda = 2$, $\lambda = 6$, $\lambda = 9$ respectively | 96 |
| 6.12 Final steady state patterns and time evolution of v | 97 |
| 6.13 μ_1 and μ_2 curves for different values of λ when $d = 0.001$ | 98 |
| 6.14 λ_1 and λ_2 curves for different values of d when $\lambda = 2.1$ | 99 |
| 6.15 Branching structures of some corals. (a) Branching coral (b) Mushroom coral. Reproduced from [13] | 100 |
| 6.16 Growth bands of two specimens of coral (a) low-density annual growth band in their apex (b) high-density annual growth band in their apex. Reproduced from [14] | 100 |
| 6.17 Sketch for the aggregation of solid particles(produced by reaction-diffusion process) on the top surface of the growing solid skeleton. (a):initial state, (b): starting state of the branching (b): two branches state | 101 |

| | |
|---|-----|
| 6.18 Sketch for the development of 3D branching structure form a set of density plots (If the density plot is multiplicative spot) of v at different time steps of the 2D model. | 102 |
| 7.1 Density plot of $v(x, t)$ for different values of α and λ . (a) for $\lambda = 0.5$, (b) for $\lambda = 2.1$, (c) for $\lambda = 10$, (d) for $\lambda = 20$. | 104 |
| 7.2 Density plot of $v(x, t)$ for $\lambda = 2.1$, $\alpha = 4.2048$ and different domain sizes | 105 |
| 7.3 Density plot of $v(x, t)$ for $\lambda = \lambda_k = 1.1 + k$, $\alpha = \alpha_k = \frac{\lambda_k^2}{\sqrt{\lambda_k - 1}}$ for $k = 1, 2, \dots, 9$. Figures in first raw for: $k = 1, 2, 3$, second raw for: $k = 4, 5, 6$ and third raw for: $k = 7, 8, 9$ from left to right in each raw | 106 |
| 7.4 Spots initiation and multiplication process for the case $\alpha = 1.0$, $\lambda = 0.5$ | 107 |
| 7.5 Spots initiation and multiplication process for the case $\alpha = 4.205$, $\lambda = 2.1$ | 108 |
| 7.6 Spots initiation and multiplication process for the case $\alpha = 9.2376$, $\lambda = 4.0$ | 108 |
| 7.7 Spots initiation and multiplication process for the case $\alpha = 40$, $\lambda = 4.0$ | 109 |
| 7.8 Isosurfaces of $v(x, y, t)$ for some parameter values which are lie on the lower boundary of the Turing space | 110 |
| 7.9 Isosurfaces of $v(x, y, t)$ for some parameter values which are deviated from the lower boundary of the Turing space | 111 |
| 7.10 Sketch for α_1 and α_2 in Turing space | 112 |
| 7.11 Patterns for linearly time dependent α . | 113 |
| 7.12 Patterns for quadratically varying α with respect to time | 114 |
| 7.13 Patterns for varying α with respect to the fourth power of time | 115 |
| 8.1 Non-existence parameter region | 121 |

List of Tables

| | | |
|-----|---|-----|
| 5.1 | Classification of the equilibrium point S_2 | 62 |
| 8.1 | Number of negative eigenvalues for $\mathbf{w}_* = \mathbf{w}_2$ when $\mu_m = 0$ | 128 |
| 9.1 | Steady states in different parameter regions | 133 |
| 9.2 | Stability of S_2 in different parameter regions | 133 |
| D.1 | Format of the program output | 162 |