

Table of Contents

Declaration.....	II
Table of Contents.....	III
List of Tables.....	VIII
List of Figures.....	X
List of Appendices.....	XI
Abbreviations.....	XII
Abstract.....	XIII
Acknowledgements	XV

Chapter 1: General Introduction

Background.....	2
1.1 Study species - Scalloped Spiny Lobster (<i>Panulirus homarus</i>)	
1.1.1 Species taxonomy and identification.....	2
1.1.2 Distribution.....	4
1.1.3 Ecology.....	6
1.1.4 Life cycle.....	7
1.1.5 Economical importance and lobster fishery.....	8
1.1.6 Management.....	9
1.2. Approach	
1.2.1Molecular methods.....	10
1.2.1.1 Population genetic.....	10

1.2.1.2 Molecular markers.....	13
1.2.1.2.1 Mitochondrial DNA markers.....	14
1.2.1.2.2 Microsatellites.....	17
1.2.2 Habitat characteristics.....	18
1.2.2.1 Habitat heterogeneity.....	21
1.2.2.2 Environmental parameters.....	22
1.2.2.3 Spatial habitat mapping.....	24
1.2.2.4 Geographic Information System (GIS).....	26
1.3 Thesis rationale and objectives.....	27

Chapter 2: DNA barcode of *Panulirus* species and investigation of phylogenetic status of *P. homarus* recorded from the Southern coastal region of Sri Lanka using mitochondrial DNA markers

2.1 Introduction.....	30
2.2 Materials and Methods.....	33
2.2.1 Study area.....	33
2.2.2 Sample collection and preservation.....	33
2.2.3 DNA isolation.....	34
2.2.4 DNA quantification.....	34
2.2.5 PCR amplification.....	34
2.2.6 Gel electrophoresis.....	35
2.2.7 DNA Sequencing.....	35
2.2.8 Alignment and phylogenetic analyses.....	35

2.3 Results.....	36
2.3.1 Sequence characteristics/Nucleotide Composition of COI and 16S alignments.....	36
2.3.2 Phylogenetic relationships.....	38
2.3.2.1 COI data analysis.....	38
2.3.2.2 16S rRNA data analysis.....	39
2.4 Discussion.....	52
2.4.1 Phylogenetic relationship.....	52
2.4.2 DNA barcoding.....	53
2.4.3 Evolutionary history.....	54
2.5 Conclusion.....	55

Chapter 3: Population genetic structure of Scalloped Spiny Lobster, *Panulirus homarus* in the Southern Coast of Sri Lanka inferred from mitochondrial DNA sequences and microsatellite DNA data

3.1 Introduction.....	57
3.2. Materials and Methods.....	60
3.2.1 Collection of Samples.....	60
3.2.2 Genomic DNA extraction.....	60
3.2.3 MtDNA Primer design, Amplification and Sequencing.....	61
3.2.4 MtDNA sequences analysis.....	62
3.2.5 Microsatellite amplification.....	63
3.2.6 Microsatellite band scoring.....	64
3.2.7 Microsatellite Data analysis.....	64

3.2.7.1 Genetic diversity.....	64
3.2.7.2 Population structure.....	64
3.3 Results.....	65
3.3.1 MtDNA analysis.....	65
3.3.1.1 Genetic variation.....	65
3.3.1.2 Population genetic structure.....	71
3.3.1.3 Neutrality estimates and <i>P. homarus</i> demographic history.....	73
3.3.2 Microsatellite analysis.....	75
3.3.2.1 PCR amplification.....	75
3.3.2.2 Genetic variability in microsatellite loci.....	75
3.3.2.3 Population genetic structure.....	78
3.4 Discussion.....	79
3.4.1 Genetic diversity.....	79
3.4.2 Population genetic structure.....	82
3.5 Conclusion.....	85


Chapter 4: Habitat characteristic analysis of scalloped spiny lobster (*Panulirus homarus*) in Southern coastal region of Sri Lanka using GIS technology

4.1 Introduction.....	87
4.2 Materials and Methods.....	89
4.2.1 Study area.....	89
4.2.2 Sampling design.....	90
4.2.3 Water quality parameters.....	93

4.2.4 Spatial analysis.....	93
4.3 Results.....	93
4.4 Discussion.....	100
4.5 Conclusion.....	103

Chapter 5: General conclusions, Recommendations for Lobster fishery management and Future researches

5.1 General Conclusions.....	106
5.2 Recommendations for lobster fishery management.....	109
5.3 Future researches.....	111
References.....	113



List of Tables

Table 2.1: Locality, BOLD accession and NCBI identification of Sri Lankan samples which used in COI data analysis

Table 2.2: Locality, Genbank accession and NCBI identification of Sri Lankan samples which used in 16S rRNA data analysis

Table 2.3: Locality information and GenBank accession numbers of the COI sequences downloaded from the NCBI GenBank database

Table 2.4: Locality information and GenBank accession numbers of the 16S rRNA sequences downloaded from the NCBI GenBank database

Table 2.5: Percentage of sequence divergence levels for 16S rRNA gene region for *Panulirus* species used from the SCSL

Table 2.6: Percentage of sequence divergence levels for COI gene region for *Panulirus* species used from the SCSL

Table 3.1: Generated primer sequences for mtDNA CytB and D-Loop gene amplification for *P. homarus*

Table 3.2: Primers and PCR conditions for the amplification of mtDNA gene regions

Table 3.3: List of DNA microsatellites and cross amplified primer-pairs for *P. homarus*

Table 3.4: Results of molecular diversity estimation

Table 3.5: Location of the SCSL where *P. homarus* were sampled, number of individuals per sampling site (N_a), distributions of haplotypes and summary of genetic variability statistics; number of haplotypes (N_h), haplotype diversity (H_d), nucleotide diversity (N_d) and number of polymorphic sites (N_p) at three mtDNA loci.

Table 3.6: AMOVA results for *Panulirus homarus* among four locations from the southern coast of Sri Lanka

Table 3.7: Pairwise fixation index (F_{ST} values) between *Panulirus homarus* populations (based on D-Loop, COI and CytB gene regions)

Table 3.8: Neutrality indexes and Mismatch analyses for *P. homarus* populations based on COI, D-Loop and CytB gene regions

Table 3.9: Genetic variability of four microsatellite loci in 04 populations for *P. homarus* in the SCSL

Table 3.10: Summary of F-Statistics and gene flow for all loci derived from microsatellite analyses

Table 3.11: Genetic variability among samples at four microsatellite loci

Table 3.12: Nei's unbiased measures of genetic identity and genetic distance

Table 3.13: Analysis of molecular variance within and among four populations of *Panulirus homarus*

Table 4.1: The geographic distribution, lobster abundance and average environmental parameters of the selected sites

List of Figures

- Figure 1.1: Characteristic features of *P. homarus* found in southern coastal region of Sri Lanka; Carapace and leg characteristics (A), head mark and front horns (B), dorsal abdominal characteristics (C) and ventral thoracic characteristics (D)
- Figure 1.2: The distribution of *P. homarus* in the Indo-west Pacific region (Holthuis, 1991)
- Figure 1.3: Schematic presentation of development of life stages of spiny lobster at different habitats (Lachlan, *et al.*, 2007)
- Figure 2.1: Study area showing localities at which *Panulirus homarus* was collected in the SCSL
- Figure 2.2: Amplified PCR products of COI gene region (indicated by arrow) of five different *Panulirus* species; L1 – *P. homarus*, L2 – *P. ornatus*, L3 – *P. longipes*, L4 – *P. versicolor*, L5 – *P. penicillatus* and M – DNA marker
- Figure 2.3: Amplified PCR products of 16S rRNA gene region (indicated by arrow) of four different *Panulirus* species; L1 – *P. homarus*, L2 – *P. ornatus*, L3 – *P. versicolor*, L4 – *P. penicillatus* and M –DNA marker
- Figure 2.4: The genus-specific ML tree derived from analyses of COI gene region of *Panulirus* species
- Figure 2.5: The genus-specific NJ tree derived from analyses of COI gene region of *Panulirus* species
- Figure 2.6: The genus-specific MP tree derived from analyses of COI gene region of *Panulirus* species
- Figure2.7: The genus-specific ML tree derived from analyses of 16S rRNA gene region of *Panulirus* species
- Figure2.8: The genus-specific NJ tree derived from analyses of 16S rRNA gene region of *Panulirus* species
- Figure2.9: The genus-specific MP tree derived from analyses of 16S rRNA gene region of *Panulirus* species

Figure 3.1: Sampling sites of four different populations of spiny lobsters in Southern coast of Sri Lanka

Figure 3.2: Haplotype networks for mtDNA COI gene region

Figure 3.3: Haplotype networks for mtDNA CytB gene region

Figure 3.4: Dendrogram Based Nei's (1978) Genetic distance: Method = UPGMA; Modified from NEIGHBOR procedure of PHYLIP Version 3.5

Figure 4.1: Sampling habitat areas of *P. homarus* in the Southern coast of Sri Lanka

Figure 4.2: Google images of the selected locations and sub sampling sites

Figure 4.3: Spatial distribution of Environmental parameters of Godawaya site

Figure 4.4: Spatial distribution of Environmental parameters of Hikkaduwa site

Figure 4.5: Spatial distribution of Environmental parameters of Weligama site

Figure 4.6: Spatial distribution of Environmental parameters of Patanagalle site

Figure 4.7: Percentage of bottom substrate conditions of the four selected sites

List of Appendices

Appendix I: The water current pattern of the North Indian Ocean

Appendix II: Related publications of chapter 2

Appendix III: Related publications of chapter 3