

## TABLE OF CONTENTS

CHAPTER	PAGE
I. INTRODUCTION . . . . .	1
II. REVIEW OF LITERATURE . . . . .	4
A. Relations of -SH groups to frost hardiness and growth . . . . .	4
B. -SH groups in proteins, peptides, and amino acids . . . . .	8
C. Methods used to measure -SH groups . . .	19
1. Oxidizing agents . . . . .	19
2. Nitroprusside . . . . .	25
3. Alkylating agents . . . . .	27
4. Mercaptide forming agents . . . . .	28
a. Spectrophotometric . . . . .	28
b. Polarographic . . . . .	32
c. Amperometric . . . . .	33
5. Other methods . . . . .	41
6. Complications due to oxidation and denaturation . . . . .	43
7. Disulfide reduction . . . . .	52
III. MATERIALS AND METHODS . . . . .	59
A. Determinations of -SH groups . . . . .	59

CHAPTER	PAGE
B. Preparation of plant materials . . . . .	86
IV. RESULTS . . . . .	91
A. Amperometric measurements of known compounds . . . . .	91
1. Current-voltage curves . . . . .	91
2. n-Dodecylmercaptan . . . . .	98
3. Cysteine . . . . .	103
4. Glutathione . . . . .	106
5. Thiogel . . . . .	110
6. Cupric titrations . . . . .	114
7. Antifoaming agent . . . . .	115
B. Measurements of -SH groups of plant homogenates . . . . .	120
C. Refinements of the amperometric methods .	180
1. The use of alcohol and sulfite reduction . . . . .	180
2. Mercuric titrations vs silver titrations . . . . .	192
V. DISCUSSION AND CONCLUSIONS . . . . .	213
VI. SUMMARY . . . . .	225
BIBLIOGRAPHY . . . . .	229

## LIST OF TABLES

TABLE	PAGE
1. Titrations of -SH groups of n-dodecylmercaptan in ammonia buffer, in air and under nitrogen .	101
2. Titrations of -SH groups of cysteine in ammonia buffer, in air and under nitrogen. . .	105
3. Titrations of -SH groups of glutathione in ammonia buffer, in air and under nitrogen. . .	107
4. Titrations of -SH groups of oxidized gluta- thione before and after reduction with different reducing agents. . . . .	108
5. The effect of time on aqueous solutions of GSH, titrated with silver nitrate, under nitrogen, in 13% methanol. . . . .	110
6. Titrations of -SH groups of thiogel in ammonia buffer, in air and under nitrogen. . . . .	112
7. Cupric titrations of cysteine and plant fractions. . . . .	116
8. The use of octyl alcohol as an anti-foaming agent when making titrations of -SH groups . .	118
9. The use of octyl alcohol in recovering the -SH groups from the foam of soybean homogenates. .	119

TABLE	PAGE
10. Changes in -SH groups with changes in frost resistance of <u>Saxifraga stracheyi</u> . . . . .	132
11. Changes in -SH groups of <u>Saxifraga stracheyi</u> leaves incapable of frost hardening . . . . .	133
12. Changes in -SH groups with changes in frost resistance of cabbage . . . . .	134
13. Changes in -SH groups of <u>Pinus sylvestris</u> needles with changes in frost resistance. . .	137
14. Changes in -SH groups of tender plants incapable of hardening when exposed to low temperature . . . . .	138
15. The -SH groups of fifteen wheat varieties, ranked according to field proven frost hardi- ness, as determined on the sprouts of seed- lings in the coleoptile to one-leaf stage . .	139
16. A comparison between the -SH groups found in homogenates of wheat varieties in which the titrations were made under nitrogen, in air, and after standing 24 hours . . . . .	140
17. A comparison of the -SH groups found in super- natants of thirteen wheat varieties when	

TABLE	PAGE
determined immediately after preparation and after standing 24 hours at + 3°C . . . . .	142
18. A comparison between -SH groups of wheat leaves sampled in the morning and in late afternoon . . . . .	144
19. A comparison between two 15-second and two 30-second blendings of wheat leaves and -SH groups of homogenates, plastids, and super- natant fractions . . . . .	145
20. Changes in -SH groups of homogenate, plastid supernatant, grana, and grana supernatant fractions with changes in frost resistance of <u>Saxifraga stracheyi</u> . Leaves homogenized in nitrogen . . . . .	147
21. Changes in -SH groups of homogenate, plastid, plastid supernatant, and grana supernatant with changes in frost resistance of <u>Saxi- fraga stracheyi</u> . Leaves homogenized in air .	149
22. Changes in -SH groups of homogenates and mitochondria with changes in frost resist- ance of <u>Saxifraga stracheyi</u> leaves. . . . .	151



TABLE	PAGE
23. Changes in -SH groups of <u>Saxifraga stracheyi</u> leaf homogenates blended in air and under nitrogen and disulfides reduced with sodium sulfite giving the total possible -SH groups. . . . .	152
24. Changes in -SH groups of cabbage homogenates, plastid, and plastid supernatant fractions with exposure to frost hardening conditions .	154
25. GSH oxidizing activity of <u>Saxifraga stracheyi</u> blended in nitrogen or in air . . . . .	158
26. GSH oxidizing activity of cabbage blended in nitrogen or in air. . . . .	159
27. GSH oxidizing activity of Scotch pine blended in nitrogen or in air . . . . .	161
28. GSH oxidizing activity of sunflower and kidney bean blended in nitrogen or in air. . . . .	162
29. Changes in -SH groups of <u>Saxifraga stracheyi</u> homogenates when exposed to hardening temperatures, blended in nitrogen or in air, and reduced with sodium sulfite in the absence or presence of guanidine hydrochloride	164

TABLE	PAGE
30. Changes in -SH groups of sunflower homogenates when exposed to hardening temperatures, blended in nitrogen or in air, and reduced with sodium sulfite in the presence of guanidine hydrochloride, and bean homogenates only blended in nitrogen or in air. . . . .	165
31. The effects of increased concentration of sodium sulfite on reduction of -SS- bonds of <u>Saxifraga stracheyi</u> homogenates . . . . .	167
32. The -SH groups of <u>Saxifraga stracheyi</u> homogenates in the native state, reduced with sodium sulfite, denatured with guanidine hydrochloride, and reduced with sodium sulfite in the presence of guanidine hydrochloride . . . . .	169
33. A comparison between the -SH groups of cabbage homogenates titrated in aqueous media and in 13% alcohol . . . . .	172
34. The -SH groups of <u>Saxifraga stracheyi</u> homogenates as determined by the regular titration method and with excess silver titrant added	

TABLE	PAGE
before the homogenates were added to the titration medium . . . . .	175
35. Differences in the -SH groups of cabbage homogenized at pH 7.5 or pH 4.5 . . . . .	177
36. Changes in -SH groups of <u>Saxifraga stracheyi</u> and cabbage leaves when frozen above and below the killing temperature . . . . .	179
37. A comparison of the -SH groups of thiogel titrated with mercuric chloride at the RMPE and silver nitrate at the RPWE. . . . .	208



## LIST OF FIGURES

FIGURE	PAGE
1. Reference and indicator electrodes used in amperometric titrations . . . . .	66
2. Apparatus for amperometric titrations employing a non-synchronous motor, microammeter, and cell with fixed potential of -0.23V . . . . .	69
3. Amperometric titration of <u>Saxifraga stracheyi</u> homogenate showing the relation of current at a fixed-point to the plotted end-point . . . . .	71
4. Amperometric titration of cabbage homogenate and thiogel showing the relation of current at a fixed-point to the plotted end-point . . . . .	72
5. Diagram of apparatus for manual control of applied voltages for current-voltage curves and amperometric titrations . . . . .	77
6. Amperometric titration vessel equipped with nitrogen flow to insure oxygen free medium . . . . .	78
7. Amperometric titration of <u>Saxifraga stracheyi</u> homogenate in aqueous ammonia buffer and in 13% methanol . . . . .	83
8. Current-voltage curves for n-dodecylmercaptan	

FIGURE	PAGE
in aqueous ammonia buffer or 95% alcohol, with and without silver nitrate added . . . .	92
9. Current-voltage curves for cysteine in aqueous ammonia buffer with and without silver nitrate added . . . . .	94
10. Current-voltage curves for cysteine in ammonia buffer in 95% alcohol with and without silver nitrate added . . . . .	95
11. Current-voltage curves for reduced glutathione in aqueous ammonia buffer with and without silver nitrate added . . . . .	96
12. Current-voltage curves for reduced glutathione in ammonia buffer in 95% alcohol with and without silver nitrate added . . . . .	97
13. Amperometric titration of n-dodecylmercaptan in aqueous ammonia buffer or in ammonia buffer in absolute ethanol . . . . .	99
14. Current-voltage curves for thiogel in ammonia buffer . . . . .	111
15. Current-voltage curves for ammonia buffer only, ammonia buffer made with 0.166M sucrose and	

FIGURE	PAGE
0.0133M $K_2HPO_4$ , and with silver nitrate added . . . . .	122
16. Blank amperometric titration of ammonia buffer, pH 9.3, in the presence of 0.166M sucrose and 0.0133M $K_2HPO_4$ . . . . .	123
17. Current-voltage response for several additions of silver nitrate in ammonia buffer in the presence of 0.166M sucrose and 0.0133M $K_2HPO_4$ .	123
18. Current-voltage curves for Seneca wheat in aqueous ammonia buffer in the presence of sucrose and phosphate, with and without silver nitrate added . . . . .	124
19. Current-voltage curves for <u>Saxifraga stracheyi</u> homogenate in aqueous ammonia buffer in the presence of sucrose and phosphate, with and without silver nitrate added . . . . .	125
20. Current-voltage curves for cabbage homogenate in aqueous ammonia buffer in the presence of sucrose and phosphate, with and without silver nitrate added . . . . .	126
21. Current-voltage curves for sunflower homogenate	

FIGURE	PAGE
in aqueous ammonia buffer in the presence of sucrose and phosphate, with and without silver nitrate added . . . . .	127
22. Current-voltage curves for aqueous ammonia buffer with sodium sulfite added and with sodium sulfite plus silver nitrate added . . .	129
23. Current-voltage curves for ammonia buffer in 60% ethanol with sodium sulfite added and with sodium sulfite plus silver nitrate added.	130
24. Amperometric titration of <u>Saxifraga stracheyi</u> homogenate with excess silver nitrate added to the supporting medium before the homogenate was added and then extrapolated to the end-point . . . . .	174
25. Amperometric titration of supernatant from homogenized <u>Saxifraga stracheyi</u> leaves in ammonia buffer and in ammonia buffer with sodium sulfite added . . . . .	183
26. Amperometric titration of cysteine in aqueous ammonia buffer with and without sodium sulfite added . . . . .	183

FIGURE	PAGE
27. Amperometric titration of cysteine in ammonia buffer in 89% ethanol with and without sodium sulfite added . . . . .	184
28. Amperometric titration of reduced glutathione in aqueous ammonia buffer with and without sodium sulfite added . . . . .	184
29. Per cent recovery of cysteine in increasing concentrations of ethanol by argentometric amperometric titrations at the RPWE in ammonia buffer, (a) without sodium sulfite added and (b) with sodium sulfite added . . .	185
30. Per cent recovery of reduced glutathione in increasing concentrations of ethanol by argentometric amperometric titrations at the RPWE in ammonia buffer, (a) without sodium sulfite added and (b) with sodium sulfite added . . . . .	186
31. Amperometric titration of cysteine in ammonia buffer in 60% ethanol with and without sodium sulfite added . . . . .	189
32. Amperometric titration of supernatant from	

FIGURE	PAGE
homogenized cabbage leaves in ammonia buffer in 60% ethanol with and without sodium sul- fite added . . . . .	189
33. Amperometric titration of a mixture of cysteine and reduced glutathione in ammonia buffer in 60% ethanol . . . . .	190
34. Current-voltage curves for tris buffer only, tris buffer plus 0.01M NaCl, and tris buffer plus 0.10M NaCl, at the RMPE . . . . .	195
35. Current-voltage curves for tris buffer at the RMPE with and without excess mercuric chloride added. All with 0.1M NaCl added. . .	196
36. Current-voltage curves for cysteine in tris buffer at the RMPE with and without mercuric chloride added . . . . .	197
37. Amperometric titration of cysteine in tris buffer at the RMPE with mercuric chloride as titrant . . . . .	198
38. Current-voltage curves for reduced glutathione in tris buffer at the RMPE with and without excess mercuric chloride added . . . . .	199



FIGURE	PAGE
39. Amperometric titration of reduced glutathione in tris buffer and 0.01M NaCl at the RMPE with mercuric chloride as titrant showing double end-points . . . . .	201
40. Amperometric titration of reduced glutathione in tris buffer and 0.05M NaCl at the RMPE with mercuric chloride as titrant showing double end-points . . . . .	202
41. Current-voltage curve for thiogel in tris buffer and 0.01M NaCl at the RMPE. . . . .	204
42. Amperometric mercuric titrations of thiogel in tris buffer at the RMPE (a) at pH 7.5 and (b) at pH 1.95. Both titrations in the presence of 0.10M NaCl . . . . .	205
43. Amperometric mercuric titrations of thiogel in tris buffer at the RPWE (a) at pH 7.4, (b) pH 2.9 and (c) in phosphate buffer at pH 8.3 . . . . .	206
44. Amperometric argentometric titrations of thiogel at the RPWE in ammonia buffer, pH 8.6 and in tris buffer, pH 7.4 . . . . .	207

FIGURE	PAGE
45. Amperometric argentometric titrations of thiogel in ammonia buffer in the presence of 0.10M sodium sulfite . . . . .	211
46. Current-voltage curves at the RPWE for (a) ammonia buffer, (b) ammonia buffer with thiogel added, (c) ammonia buffer with sodium sulfite added, and (d) ammonia buffer with thiogel, sodium sulfite and excess silver nitrate added. . . . .	212