10

CONTENTS

	Page	
ACKNOWLEDGEMENTS		
TABLE OF CONTENTS		
LIST OF FIGURES		
LIST OF TABLES		
ABSTRACT		
1.0 INTRODUCTION	1	
2.0 MATERIALS AND METHODS	9	
2.1 Dietary Ingredients and Diets	9	
2.2 Preparation of Diets	12	
2.3 Experimental Setup	14	
2.3.1 Fish Size	16	
2.3.2 Faeces	17	
2.3.3 Carcass Analysis	18	
2.3.4 Experimental Conditions	18	
2.4 Chemical Analysis	18	
2.5 Statistical Analysis	19	
3.0 RESULTS	20	
3.1 Physical Conditions	20	
3.2 Mortality	20	
3.3 Behavioural Observations	21	
3.4 Food Intake	21	
3.5 Digestibility	33	

	3.6 Growth		37
	3.6.a	Average Daily Gain	41
	3.6.6	Food Conversion Ratio	44
	3.6.c	Protein Efficiency Ratio	47
	3.7 Carcass	Composition	45
	3.7.a	Protein Deposition	50
	3.7.b	Lipid Deposition	50
	3.7.0	Ash Deposition	54
	3.7.d	Energy Content	54
	3.7.e	Moisture Content	54
	3.7.f	Net Protein Utilization	54
4.0	DISCUSSION		58
5.0	CONCLUSION		74
6.0	REFERENCES		76

LIST OF FIGURES

Figure	<u>Title</u>	Fage
1	Examples to show the daily variation in food consumption expressed as % body weight per day in Ω . niloticus (L.) fry maintained on different diets.	23
2	The relationship of the overall mean intake per week expressed as (a) per g fish per day to mean body weight for diets at the three protein levels of 20%, 25% and 30%.	26
3(a)	The relationship of the protein intake per fish per day to mean body weight for diets at the three protein levels of 20%, 25% and 30%.	31
उ(b)	The relationship of the energy intake (kj) per fish per day to mean body weight for diets at the three protein levels of 20%, 25% and 30%.	32
4	The dry matter, protein and lipid digestibility for the two size groups of O. niloticus (L.) maintained on the diets of three protein levels (20%, 25% and 30%) in relation to the substitution of green gram.	36
5 	Fortnightly mean weight of $\underline{0}$. niloticus (L.) of size group 2 maintained under different dietary regimes.	39
6	Fortnightly mean weight of $\underline{0}$. <u>niloticus</u> (L.) of size group 1 maintained under different dietary regimes.	40
7	The %ADG for the two size groups of $\underline{0}$. niloticus maintained on diets of three protein levels in relation to the degree of substitution of green gram.	43

8	The relationship of FCR to the degree of green gram substitution level at three protein levels.	46
9	PER of the two size groups of <u>O. niloticus</u> fry maintained on diets of three protein levels in relation to the degree of substitution of green gram.	48
10	The changes in carcass protein, lipid and ash of <u>O</u> . <u>niloticus</u> of the two size groups maintained on different dietary regimes.	52
11	The changes in carcass energy, moisture and NPU of $\underline{0}$. <u>niloticus</u> of the two size groups maintained on different dietary regimes.	53
12	The NPU for the two size groups of <u>O</u> . <u>niloticus</u> fed on diets of three protein levels in relation to the degree of green	56

LIST OF TABLES

<u>Table</u>	<u>Title</u>	<u>F'age</u>
1(a)	The proximate composition (dry weight) and amino acid composition of \underline{F} . aureus.	10
1(b)	The proximate composition (dry weight) of fishmeal, soybean, coconut meal and wheat flour.	10
2	Composition of the mineral and vitamin mixtures.	11
3	Summary of the experimental design.	12
4	The ingredient and proximate composition of the test diets.	13
5	The values of the exponent and the constant of the statistical relationships of the weight of food consumed per g fish per day for the experimental diets and the overall intake for each group of diets, in relation to mean body weight.	25
6	The statistical relationships of the weight of food consumed per fish per day for all experimental diets and the overall intake for each group of diets, in relation to mean body weight.	29
7	Statistical relationships of the protein and energy intake per fish per day in relation to the mean body weight at three different protein levels in <u>O</u> . <u>niloticus</u> .	30
8	The percent drymatter, protein and lipid digestibility of all the diets fed to $\underline{0}$. niloticus young of two size groups.	35

9	The final mean weight and standard deviation of each group of $\underline{0}$, <u>niloticus</u> maintained under different diets.	<u>3</u> 8
10	The %ADG and FCR of <u>O</u> . <u>niloticus</u> young maintained on different dietary regimes.	42
11	The PER of $\underline{0}$. <u>niloticus</u> maintained on different dietary regimes.	47
12	Carcass composition; protein $\%$; lipid $\%$; ash $\%$; energy (kj/g); moisture $\%$; of \underline{O} . niloticus maintained on different dietary regimes.	51
13	The %NPU of $\underline{0}$. $\underline{\text{niloticus}}$ maintained on different dietary regimes.	55
14	A summary of the plant protein sources used in tilapia feeds.	60

•