

C O N T E N T S

	Page
ACKNOWLEDGEMENTS	iii
TABLE OF CONTENTS	v
LIST OF FIGURES	viii
LIST OF TABLES	x
LIST OF APPENDICES	xii
ABSTRACT	xiii
INTRODUCTION	1
PART A DETERMINATION OF OPTIMUM DIETARY PROTEIN LEVEL AND SALINITY FOR <u>Oreochromis niloticus</u> (L.) FRY.	
CHAPTER I STUDIES ON THE EFFECTS OF DIETARY PROTEIN LEVEL AND SALINITY ON PERFORMANCE OF <u>Oreochromis niloticus</u> (L.) FRY.	
I.0 INTRODUCTION	9
I.1 MATERIALS AND METHODS	10
I.2 RESULTS	17
I.2.1 Some Observed Biological and Physical Parameters	17
I.2.2 Growth	21
a. Average Daily Gain	23
b. Percentage Weight Gain	23
c. Specific Growth Rate	23
I.2.3 Food Conversion Ratio	27
I.2.4 Protein Efficiency Ratio	27
I.3 DISCUSSION	29

CHAPTER 2	EFFECTS OF DIETARY PROTEIN LEVEL AND SALINITY ON DIGESTIBILITY WITH OBSERVATIONS ON VARIABILITY IN DAILY DIGESTIBILITY.	
	2.0 INTRODUCTION	35
	2.1 MATERIALS AND METHODS	37
	2.2 RESULTS	39
	a. Digestibility in Relation to Dietary Protein Level and Salinity.	39
	b. Variability in Daily Digestibility	45
	2.3 DISCUSSION	45
CHAPTER 3	STUDIES ON CHEMICAL COMPOSITION OF <u>Oreochromis niloticus</u> (L.) FRY IN RELATION TO THE DIETARY PROTEIN LEVEL AND SALINITY.	
	3.0 INTRODUCTION	52
	3.1 MATERIALS AND METHODS	53
	3.2 RESULTS	55
	3.3 DISCUSSION	61
PART B	DETERMINATION OF THE OPTIMAL RATION SIZE AND FEEDING FREQUENCY FOR THE OPTIMAL GROWTH OF <u>Oreochromis niloticus</u> (L.) FRY.	
CHAPTER 4	DETERMINATION OF THE EFFECT OF OPTIMAL RATION SIZE AND FEEDING FREQUENCY ON THE PERFORMANCE OF <u>Oreochromis niloticus</u> (L.) FRY.	
	4.0 INTRODUCTION	72
	4.1 MATERIALS AND METHODS	73
	4.2 RESULTS	74
	4.3 DISCUSSION	86

GENERAL DISCUSSION	92
LITERATURE CITED	103
APPENDIX I	121
APPENDIX II	124

LIST OF FIGURES

<u>Figure</u>	<u>Title</u>	<u>Page</u>
1.	A typical mortality curve observed for <u>O.niloticus</u> (L.) at 15‰ salinity.	19
2.	Fortnightly changes in the mean weight of young <u>Oreochromis niloticus</u> (L.) reared on seven diets (Table 1) at four experimental salinities.	22
3.	Change in average daily gain (ADG), percentage weight gain (% day ⁻¹), food conversion ratio (FCR) and protein efficiency ratio (PER) of young <u>Oreochromis niloticus</u> (L.) in relation to dietary protein level at four experimental salinities.	24
4.	Relationship of the overall mean percent dry matter digestibility of the different diets to the dietary protein content at the four experimental salinities.	42
5.	Relationship of the overall mean percent apparent protein digestibility of the experimental diets to the dietary protein content of <u>Oreochromis niloticus</u> (L.) fry reared in fresh water, 5‰, 10‰, and 15‰ salinities.	43
6.	The daily variability of the dry matter and protein digestibility of <u>O.niloticus</u> (L.) fry maintained on diets 4 and 5 at 5‰ and 10‰ salinities.	46
7.	Changes in the proximate composition of the carcass of <u>O.niloticus</u> (L.) fry maintained on different dietary levels and environmental salinities. Percentage ash, percentage protein, percentage lipid and energy (KJg ⁻¹) are expressed on a dry weight basis.	57
8.	Relationship of the percent protein retained (expressed as percentage of dry weight) to different dietary protein level in <u>Oreochromis niloticus</u> (L.) fry.	63

9. Relationship of the percent protein retained (expressed as a percentage of dry weight) to environmental salinity in Oreochromis niloticus(L.) fry. 64
10. Fortnightly changes in the mean weight of young Oreochromis niloticus(L.) on different rations, fed twice a day. 78
11. Fortnightly changes in the mean weight of young Oreochromis niloticus(L.) fed ad libitum but at different frequencies. 79
12. Changes in percentage weight gain ($\% \text{ day}^{-1}$) food conversion ratio (FCR) and protein efficiency ratio (PER) of young Oreochromis niloticus(L.) fry reared on different rations. 82
13. Changes in percentage weight gain ($\% \text{ day}^{-1}$) food conversion ratio (FCR) and protein efficiency ratio (PER) of young Oreochromis niloticus(L.) fry reared at different feeding frequencies. 83
14. Relationship of the overall mean percent dry matter and mean percent apparent protein digestibility in Oreochromis niloticus(L.) fry maintained on different rations. 87
15. Relationship of the overall mean percent dry matter and mean percent apparent protein digestibility to the feeding frequencies in Oreochromis niloticus(L.) fry. 88

LIST OF TABLES

<u>Table</u>	<u>Title</u>	<u>Page</u>
1.	The percentage ingredients of the test diets and their proximate composition.	15
2.	The mean, range and the S.D. in temperature, pH and dissolved oxygen in each experimental 'system' during the morning and the afternoon.	20
3.	The mean final weight (\pm SE), average daily weight gain (ADG) and the growth of <u>O.niloticus</u> (L.) fry, maintained on the seven experimental diets at four salinities.	25
4.	The food conversion ratio (FCR) and protein efficiency ratio (PER) of <u>O.niloticus</u> (L.) fry maintained on experimental diets at different salinities.	28
5.	The protein levels producing maximum growth of tilapias.	30
6.	The overall mean percent dry matter and apparent protein digestibility in <u>O.niloticus</u> (L.) fry maintained on the seven experimental diets at four salinities.	41
7.	Percent dry matter, protein and energy digestibility in <u>O.niloticus</u> (L.) fry maintained on the four experimental diets at three salinities.	44
8.	Data pertaining to the extent of daily variability in percent dry matter and apparent protein digestibility of <u>O.niloticus</u> (L.) fry at salinities of 5% and 10% maintained on Diet 4 and 5.	47
9.	Changes in the proximate composition of the carcass of <u>O.niloticus</u> (L.) fry maintained on different dietary levels and environmental salinities.	56

10. The percent protein retained by O.niloticus(L.) fry maintained on different dietary protein levels at the four experimental salinities. 62
11. Detail information on feeding frequencies and feeding times. 75
12. Quantity of food consumed by O.niloticus(L.) fry expressed as a percentage of the body weight per day at each feeding frequency. 77
13. Performance of O.niloticus(L.) fry as indicated by the final weight (\pm SE), ADG, percentage weight gain ($\% \text{ day}^{-1}$), FCR and PER maintained on specific ration and fed ad libitum at different frequencies. 80
14. The mean (\pm SD) percent dry matter and apparent protein digestibility in O.niloticus(L.) fry maintained on the different rations and feeding frequencies. 85

LIST OF APPENDICES

<u>Appendix</u>	<u>Title</u>	<u>Page</u>
1	Detailed methods of analysis	121
11	Influence of dietary protein and salinity levels on protein intake of <u>O.niloticus</u> (L.) fry.	124