

RESEARCH ARTICLE

EVALUATION OF PHYTOTOXICITY OF COMPOSTS PRODUCED FROM SPENT MUSHROOM SUBSTRATE USING SEED GERMINATION BIOASSAY OF *Raphanus sativus* L.

Amarasinghe SR^{1*} and Jayaweera WMCS²

¹Department of Soil Science, Faculty of Agriculture, University of Ruhuna, Kamburupitiya, 81100, Sri Lanka.

²Department of Biosystem Technology, Faculty of Technology, University of Ruhuna, Kamburupitiya, 81100, Sri Lanka.

Received: 08 July 2022, Accepted: 23 October 2022

ABSTRACT

Spent mushroom substrate (SMS) has the potential to produce compost with different additives. However, some substances and by-products produced during the process can be toxic to seed germination and plant growth. Evaluation of the phytotoxicity is important for guaranteeing compost quality. Therefore, this study was conducted to evaluate the phytotoxicity of spent mushroom substrate compost (SMSC) using a seed germination bioassay. Compost aqueous extracts (CAEs) were prepared from SMSC produced by amending different additives (green leaves and urea) with and without an inoculum. Dilution series of 25%, 50%, and 100% were tested against seed germination of radish (*Raphanus sativus* L). Germination index (GI) in all 25% CAEs of this study were phytotoxic free and they indicated the availability of phytonutrients and/or phyto-stimulants except the non-inoculated SMSC without amendments. Similarly, all 50% CAEs were phytotoxic free except *Tithonia* amended inoculated SMSC with less phytotoxicity. Of 100% CAEs, non-inoculated *Pueraria* amended SMSC was highly phytotoxic, non-inoculated *Tithonia* and urea amended SMSC were phytotoxic free, and the rest were less phytotoxic. Seed germination of *R. sativus* was not inhibited by any of the CAEs of *Tithonia* amended SMSC indicating the higher usability as an amendment in SMS. The water extractable NH_4^+ concentration was significantly higher ($p < 0.05$) in the inoculated SMSC amended with urea and it was significantly lower ($p < 0.05$) in the non-inoculated SMSC amended with *Tithonia* and inoculated SMSC without amendments. Further, the inoculated SMSC extracts showed higher performance in GI than non-inoculated SMSC. GI showed a higher negative significant correlation ($r = -0.841$, $p < 0.01$; $r = -0.778$, $p < 0.01$) with EC in both inoculated and non-inoculated compost samples, respectively. Moreover, the EC was significantly correlated with all extractable tested ions except Cu^{2+} and Ni^{2+} in inoculated treatments. pH values in all aqueous extracts were recorded in the range of 6.27 to 8.08. Extractable elements Cu^{2+} ($r = 0.645$, $p = 0.01$) and Zn^{2+} ($r = 0.577$, $p = 0.05$) had positive correlation with pH. SMS can be recycled as compost integrating nitrogenous additives and inoculum to use safely as a soil amendment.

Keywords: Compost, Phytotoxicity, Seed germination, Spent mushroom substrate

INTRODUCTION

Edible mushroom cultivation is popular worldwide due to its fast-growing nature and the possibility to obtain higher yields within a short period. The continuous yield can be obtained for about three months. Afterwards, the spent mushroom substrate (SMS) has no

longer usable among farmers and piles up or burns into ash.

Therefore, SMS is considered as waste that is obtained from the mushroom production industry. Piling up SMS after mushroom harvesting may cause various environmental problems including water contamination and nuisance (Beyer 1996). This could be re-

*Corresponding author: rajika@soil.ruh.ac.lk

utilized to produce mushrooms or used as a direct soil amendment. The growth of unwanted microorganisms in compost produced from SMS is one of the problems in utilizing it as a soil amendment. However, the SMS can be directly used as “fresh mushroom compost” after being pasteurized under steam heat (Davis and Fidanza 2011) or by the hot composting process for at least 18 days to inactivate the unwanted fungi and weed seeds (Peters *et al.* 2000). Then the secondary pile can be formed to complete the composting process up to 90 days or less depending on the substrate. In Sri Lanka, the SMS is mostly piled up on the ground. However, these are not abundantly re-utilized in Sri Lanka as compost or any other utility.

SMS can be used to prepare compost which is used as an organic amendment. It is within the perfect range for compost production having 35% to 55% (wet weight) solids and 45% to 65% (wet weight) moisture content (Stoffella and Kahn 2001).

Compost should be free of salinity, $\text{NH}_4^+\text{-N}$, heavy metals, etc. by metabolization or immobilization at maturity ensuring free phytotoxicity (Emino and Warman, 2004). Major constraints of the SMS would be the salinity which is concentrated by added lime (CaCO_3 and $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$) in mushroom production. Therefore, a risk remains due to a high salinity problem in the final spent mushroom substrate compost (SMSC). Besides salts, SMS contain $\text{NH}_4^+\text{-N}$, heavy metals and metal pollutants, such as arsenic, chromium, nickel, and selenium depending on the initial substrate (Alvarenga *et al.* 2015) which may create phytotoxicity (Wollan *et al.* 1978; Wong *et al.* 1983). High salinity levels may affect germinating seeds and young seedlings (Kebrom *et al.* 2019) and are harmful to plant growth (Singh and Kalamdhad 2013a; 2013b). The damage may be severe to salt-sensitive plants.

The bioremediation by microorganisms can eradicate some toxicants by biological mechanisms to innocuous forms (Ojuederie and Babalola 2017). The availability of these toxicants is reduced by adjusting pH, the redox

reactions and the adsorption of pollutants (Jain and Arnepalli 2018). Redox reactions can be operated in the composting process (Bolan 2013). It helps to increase microbial population by altering pH, decreasing the heavy metals, and increasing allochthonous microbial biomass and plant nutrients (Albuquerque *et al.* 2011; Chen *et al.* 2015). Microorganisms utilize different processes such as precipitation, biosorption, enzymatic conversions of heavy metals (Ojuederie and Babalola 2017). Introducing microorganisms by inoculation may help in such processes.

Phytotoxicity assessment is mandatory as a parameter in the quality profiling of compost, before utilizing them. Even though different types of chemical analysis have been carried out by many researchers (Marambe *et al.* 1992; Ramírez *et al.* 2008; Hase and Kawamura 2012; Fuentes *et al.* 2004; Tang *et al.* 2008; Siles-Castellano *et al.* 2020) to identify potential toxicants such as organic products, $\text{NH}_4^+\text{-N}$, heavy metals, pesticide residues, many are impractical due to time intensity and expansiveness. Further, the interactive effects of these toxicants cannot be evaluated properly through chemical analysis.

Therefore, calculating the germination index (GI) is a reliable method to ascertain the phytotoxic effect of compost and it evaluates the phytotoxins which can temporarily or permanently affect the growth of a plant (Warman 1999). Thus, seed germination-related bioassays have been conducted as an effective, low cost and rapid method to determine phytotoxicity (Mazumder *et al.* 2020). Such seed germination experiments can be conducted using aqueous extracts of organic amendments and incubating the selected seed varieties (Emino and Warman 2004; Barral and Paradelo 2011). Many previous studies have followed seed germination protocols as a bioassay (Kebrom *et al.* 2019). The seed germination index has introduced by Zucconi *et al.* (1981) for cress seeds (*Lepidium sativum*, L.) by considering seed radicle length and germination percentage by comparing the control in deionized water. Later Gariglio *et al.* (2002) utilized lettuce, whereas Emino and Warman

(2004) used 14 small to large-sized seed species such as *Amaranthus* (*Amaranthus tricolor*), bean (*Phaseolus vulgaris*), carrot (*Daucus carota*), radish (*Raphanus sativus*), etc. Many researchers have evaluated GI using *Raphanus sativus* seeds which are sensitive to toxicants at the early germination cycle (<48 h) and the convenient medium-seed size (Iannotti *et al.* 1994; Gajdos 1997; Luo *et al.* 2018).

As the first step of the seed germination test, the compost aqueous extracts should be prepared according to the suitable extraction ratio. There is no universal ratio of extraction and many researchers have adopted several extraction ratios (Luo *et al.* 2018). Some researchers have implemented 1:10 (w/v) aqueous extracts (Guo *et al.* 2012; Yang *et al.* 2013; Huang *et al.* 2016; Luo *et al.* 2018). Also, Said-pullicino *et al.* (2007), Pampuro *et al.* (2010) and Cesaro *et al.* (2015) have standardized the moisture content for 85% by adding deionized water into the fresh sample after determining the dry matter content of each sample. Hence, many studies have used different extraction methods, seed types and numbers, the volume of the extract and incubation time for the test. Of these, the majority have utilized Petri dishes to test the radicle elongation (Luo *et al.* 2018).

In this study, we assumed that SMS integrated with other amendments will affect the seed germination and root elongation due to the phytotoxic nutrient elements, heavy metals and metal pollutants present and the added inoculum may accelerate the decomposition and reduce the phytotoxicity. As phytotoxicity assessment is a significant and compulsory parameter to determine the quality of SMSC, the seed germination bioassay of radish (*Raphanus sativus* L.) seeds using aqueous extracts of each final compost mixtures were tested to evaluate the possible phytotoxic effect of these compost mixtures.

MATERIALS AND METHODS

Spent mushroom substrate for compost

SMS of oyster mushroom (*Pleurotus ostreatus*) obtained from mushroom

production unit of Faculty of Agriculture, University of Ruhuna, Sri Lanka was used as the key waste to prepare compost. Green leaves i.e. *Peuraria pasioloids* leaves, *Gliricidea sepium* leaves, *Tithonia diversifolia* (wild sunflower) leaves, and inorganic nitrogen fertilizer (urea) were used as N sources. The total C%, total N% content and moisture content were analyzed in all the raw materials. The SMS was amended and mixed thoroughly with the green leaves and urea separately according to the amount of materials calculated using formula 1 (Rynk *et al.* 1992). The C/N ratio of the final compost mixture for optimum decomposition was considered 25:1.

$$R = \frac{[Q_1 C_1 (100 - M_1) + Q_2 C_2 (100 - M_2)]}{[Q_1 N_1 (100 - M_1) + Q_2 N_2 (100 - M_2)]} \quad \text{..... Eqn 1}$$

where R is the C/N ratio of final compost mixture for optimum decomposition, Q_1 is the known weight of sample (A), C_1 is the known C% of sample (A), M_1 is the known moisture% of sample (A), M_2 is the known moisture % of sample (B), N_1 is the known N% of sample (A), Q_2 is the unknown weight of sample (B) to be added, C_2 is the known C% of the sample (B) and N_2 is the known N% of sample (B).

Then, another five mixtures were prepared similarly and inoculated with an inoculum/biodynamic formulation (Amarasinghe and Gunawardhana 2020). Two treatments were prepared with SMS without amending green leaves or urea but mixing with inoculum and without inoculum, respectively. Each compost mixture of 2 kg was separately put into a black polythene bag of gauge 150 and labelled accordingly. All the treatments were replicated three times and allowed to decompose aerobically. The bags were arranged in ambient temperature and humidity conditions using a Complete Randomized design (CRD). Every compost mixture was mixed thoroughly every other day to speed the decomposition process. Moisture content was adjusted by applying water by monitoring the moisture level by field method. Mixtures were allowed to decompose for 90 days.

Preparation of inoculum

To prepare the inoculum, 500 g of cow dung, 250 ml of cattle urine, 100 g of brown sugar, 100 g of mung bean flour and one handful of topsoil from an undisturbed forest area were collected and added to a 10 L bucket. The contents were mixed well and increased volume up to 10 L by adding water. The mixture was stirred well to aerate once in two days and kept for 10 days. The inoculum was applied at a rate of 500 L/100 kg of substrate considering the bulk density of the substrate. The bulk density of coarse sawdust was 340 kg/m³.

Determination of Germination index (GI)

In the present study, a germination bioassay was performed on aqueous extracts of mature SMSC using *Raphanus sativus* seeds. *R. sativus* seeds were used due to the sensitivity to toxicants in the early germination cycle and convenient seed size. Certified *R. sativus* seeds with 75% germination were used to determine the GI. GI test was conducted according to Kebrom *et al.* (2019). CAEs from ten mature SMSC, (i) SMS + *Gliricidea sepium* leaves (ii) SMS + *Gliricidea sepium* leaves with inoculum (iii) SMS + *Tithonia diversifolia* leaves (iv) SMS + *Tithonia diversifolia* leaves with inoculum (v) SMS + *Peuraria pasioloids* leaves (vi) SMS + *Peuraria pasioloids* leaves with inoculum (vii) SMS + urea (viii) SMS + urea with inoculum (ix) SMSC without inoculum (x) SMSC with inoculum were used for the seed germination bioassay.

Sample preparation and physicochemical characterization

Matured compost mixtures were dried overnight at 105 °C in an oven. The dry matter and the moisture content were determined. Then it was standardized up to 85% moisture content by adding deionized water. By Shaking the samples on a horizontal shaker in a volumetric flask for two hours and solutions were filtered using a Whatman no. 01 filter paper. Three concentrations (25%, 50%, and 100%) of the aqueous extracts were prepared to assure the accuracy of the evaluation. pH, electrical conductivity (EC), water-extractable NH₄⁺, Fe³⁺, Cu²⁺, Ni²⁺, Cr³⁺,

Zn²⁺, Na⁺, SO₄²⁻, and Ca²⁺ concentrations of each aqueous extracts were analyzed. pH and EC were measured using a pH and EC meter (HANNA HI 83099). Flame atomic emission spectrometry was used to analyze Ca²⁺ and Na⁺ (Tandon 1995). SO₄²⁻, NH₄⁺ and heavy metals were analyzed using a photometer according to the colourimetric methods (CHEMetrics 2018). The seeds were dipped in a 10% calcium hypochlorite solution for surface sterilization (Kim *et al* 2011) for 10 mins and the seeds were rinsed with sterilized water. Five radish seeds were placed on a Petri dish (9 cm) with lined qualitative filter paper. Three replicated dishes were prepared for each compost aqueous extract in different concentrations (25%, 50%, and 100%). Five millilitres of the extract were poured into each treatment Petri dish and 5 ml of deionized water to the control petri dish. The experiment was arranged in a complete randomized design. The GI test was conducted under the dark condition at room temperature (27 °C). The lids were kept on the Petri dish to prevent evaporation. The number of seeds germinated following 72 h incubation was counted and the radicle of the seedlings were measured. According to the standard formulas, (Kebrom *et al.* 2019, Mazumder *et al.* 2020) percentages of RSG, RRG and GI were calculated (equations 2, 3 and 4).

RSG = [Number of seeds germinated in compost aqueous extract / Number of seeds germinated in control] x 100% Eqn 2

RRG = [Mean radicle length of germinated seeds in compost aqueous extract (mm) / Mean radicle length of germinated seeds in control (mm)] x 100% Eqn 3

GI = [RSG x RRG] x 100% Eqn 4

Where RSG is the relative seed germination, RRG is the relative root growth and GI is the Germination index.

The above three terms are relative terms which were compared to the control having always 100. Then, whenever values of treatment exceed the control in either germination or root elongation, the values will

exceed 100 and the GI may be higher than 100 (Emino and Warman, 2004).

Statistical Analysis

The number of germinated seeds and the average radicle length in each of the replicates were determined. The RSG, RRG, and GI of each were calculated. The results reported in this paper are the mean GI, RSG, and RRG of treatments, and the error bars are the standard error of the mean (SE). The results of physical and chemical properties were also analyzed similarly. Treatment means were tested by one-way ANOVA followed by a Duncan Multiple Range Test (DMRT) for the mean separation. A correlation analysis was conducted between GI, RSG, RRG and other physicochemical parameters of treatments. All analyses were performed using SPSS 17.0. (SPSS 2006).

RESULTS AND DISCUSSION

Germination of *Raphanus sativus* seeds in different CAEs (25%, 50%, and 100%) of different treatments and deionized water (control)

GI is the most important parameter to evaluate the phytotoxicity of mature compost as it directly examines the effect of compost on seed germination and seedling growth (Wang

et al. 2020). Table 1 shows the mean values of GI in 25%, 50%, and 100% CAEs. According to Emino and Warman (2004), compost with GI values $\leq 50\%$ is highly phytotoxic, between 50% and 80% are less phytotoxic, $\geq 80\%$ are phytotoxic free, and 0 indicates extreme phytotoxicity. The availability of phytonutrients and/or phyto-stimulants is the reason for GI higher than 100% (Emino and Warman 2004). Correspondently, all 25% CAEs of this study were phytotoxic free, and they indicated the availability of phytonutrients and/or phyto-stimulants except the non-inoculated SMSC without amendments. Similarly, all 50% CAEs were phytotoxic free except *Tithonia* amended inoculated SMSC with less phytotoxicity. Of 100% CAEs, non-inoculated *Tithonia* and urea amended SMSC were phytotoxic free, non-inoculated *Pueraria* amended SMSC was highly phytotoxic, and the rest were less phytotoxic. Emino and Warman (2004) found that 25% and 50% CAEs are not phytotoxic, presumably diluted to a non-phytotoxic concentration resulted in a $GI > 80\%$, while the dilution to 75% CAE and undiluted 100% immature CAEs resulted in a $GI \leq 20\%$ and are toxic which resemble the results of the present study.

Table 1: Means of the GI values of different CAEs

Treatment	GI (%)			
	25% CAE	50% CAE	100% CAE	
SMS+Gliricidia	non-inoculated	153 a*	105 e	68 cd
	inoculated	120 h	95 g	67 d
SMS+Tithonia	non-inoculated	102 i	86 h	87 b
	inoculated	152 b	73 i	74 c
SMS+Pueraria	non-inoculated	123 g	94 g	40 f
	inoculated	135 e	109 d	72 c
SMS+Urea	non-inoculated	127 f	92 g	82 b
	inoculated	144 e	125 b	60 e
non amended SMSC	non-inoculated	92 j	111 c	72 c
	inoculated	140 d	134 a	57 e
Control (Deionized water)		100 i	100 f	100 a

* Mean values designated by the same letter are not significantly different at $p < 0.05$ as determined by DMRT.

Means of the GI values of CAEs were compared using DMRT (Table 1) to see the significant differences among calculated GI values among treatments.

In most treatments of 25% and 50% CAEs, GI percentages were significantly ($p < 0.05$) higher than the control except for the 100% CAE. Emino and Warman (2004) indicated that GI may be higher than the control, whenever the values of treatments exceed the control in either germination or root elongation. However, within different CAEs, there were significant differences ($p < 0.05$) among different treatments (Table 1).

The significantly lowest GI value at 25% CAE was observed in the treatment of non-inoculated SMSC without amendments while the highest was in *Gliricidia* amended non-inoculated SMSC ($p < 0.05$). The GI was increased as SMSC without amendments $<$ *Tithonia* amended SMSC $<$ *Pueraria* amended SMSC $<$ urea amended SMSC $<$ *Glyricidia* amended SMSC in non-inoculated compost, while *Glyricidia* amended SMSC $<$ *Pueraria* amended SMSC $<$ SMSC without amendments $<$ Urea amended SMSC $<$ *Tithonia* amended SMSC in inoculated compost.

The GI in 50% CAE, *Tithonia* amended SMSC was significantly lower ($p < 0.05$) compared to all treatments while the highest ($p < 0.05$) was observed in inoculated SMSC without amendments. The GI was increased as *Tithonia* amended SMSC $<$ urea amended SMSC $<$ *Pueraria* amended SMSC $<$ *Gliricidia* amended SMSC $<$ SMSC without amendments in non-inoculated compost, while *Tithonia* amended SMSC $<$ *Gliricidia* amended SMSC $<$ *Pueraria* amended SMSC $<$ urea amended SMSC $<$ SMSC without amendments in inoculated compost. In both 25% and 50% CAEs, the GI was higher than the control (100%) in most treatments showing the phyto-stimulant behaviour of these compost extracts resembling the study of Siles-Castellano *et al.* (2020).

The GI in 100% CAE, *Pueraria* amended SMSC was significantly lower ($p < 0.05$) than all the other treatments and the control irrespectively to inoculated and non-inoculated composts. However, CAE of 100% showed significantly higher ($p < 0.05$) GI values in non-inoculated *Tithonia* amended SMSC and urea amended SMSC compared to the other non-inoculated treatments. The GI of non-inoculated *Pueraria* amended SMSC (40%) indicated the presence of phytotoxic substances comparable to the study conducted by Kebrom *et al.* (2019) using chicken manure, milorganite, and dairy manure as three commercial organic amendments. The GI was increased as *Pueraria* amended SMSC $<$ *Gliricidia* amended SMSC $<$ SMSC without amendments $<$ urea amended SMSC $<$ *Tithonia* amended SMSC in non-inoculated compost, while SMSC without amendments $<$ urea amended SMSC $<$ *Gliricidia* amended SMSC $<$ *Pueraria* amended SMSC $<$ *Tithonia* amended SMSC in inoculated compost.

Radicle elongation is a sensitive measure contributing to the GI (Emino and Warman 2004). To obtain the effect of inoculum in seed radicle elongation, it is important to find a relationship between GI and mean radicle length. Figure 1 shows the relationship between GI and means radicle length of *R. sativus* seeds in inoculated and non-inoculated treatments. Accordingly, the GI showed a positive linear relationship with the mean radicle length of *R. sativus* seeds. Further, it was clear that the inoculated treatments had a higher positive relationship ($r^2 = 0.9235$) with GI, compared to the non-inoculated treatments ($r^2 = 0.722$). Beneficial microorganisms may enhance the bioremediation process by reducing heavy metals and organic pollutants (Ojuederie and Babalola 2017). According to him, microorganism consortia in biofilms can be utilized to remediate heavy metals based on cell metabolism. Further, Sobarzo-Bernal *et al.* (2021) observed an enhanced radicle length by a biostimulant, which indicated the effect of microbial activity on radicle elongation.




Figure 1: Relationship between GI and mean radicle length of *R. sativus* seeds in inoculated and non-inoculated treatments

pH, EC and some elemental concentrations in different CAEs (25%, 50%, and 100%) of different treatments and deionized water (control)

pH and EC values are important in seed germination. Figure 2 shows the relationship between GI and mean EC of CAEs in inoculated and non-inoculated treatments. Accordingly, there is a linear negative relationship with the mean EC for the inoculated SMSC ($r^2=0.7064$) and for non-inoculated SMSC ($r^2=0.6009$). It is visible that in low EC concentrations, the GI in CAEs of inoculated SMSC is higher than the non-inoculated SMSC.

In the present study, pH values for all extracts were in the range of 6.27 to 8.08 (data not shown), similar to the range of typical compost (Cayuela *et al.* 2008). According to Barral and Paradelo (2011), higher pH and EC values in compost inhibit seed germination. However, salinity effects are generally negligible when the EC value is $\leq 2000 \mu\text{S}/\text{cm}$ (Hoekstra *et al.* 2002). Alkaline pH values (>9) obtained in a seed germination study conducted by Siles-Castellano *et al.* (2020) have explained the low GI values, lower than 55%.




Figure 2: Relationship with GI and mean EC of compost aqueous extracts in inoculated and non-inoculated treatments

Chemical properties and compositions of CAEs

Chemical analysis of the CAEs is needed to identify potentially phytotoxic compounds. In the present study, phytotoxic analysis was

performed only for the 100% CAE, since higher germination inhibition was observed in it. Figure 3 depicts the Na⁺ and Ca²⁺ composition in the compost aqueous extracts.




Figure 3: Na⁺ and Ca²⁺ composition in the 100% CAE (Vertical bars represent standard error (mean ± SE) and mean values designated by the same letter are not significantly different at p<0.05 as determined by DMRT)

The average Na⁺ content was significantly lower (p<0.05) in non-inoculated urea amended SMSC compared to other treatments except for non-inoculated *Gliricidia* amended SMSC, non-inoculated *Tithonia* amended SMSC and inoculated SMSC without amendment. Kebrom *et al.* (2019) have found that the level of Na⁺ in the aqueous extracts was related to their EC values which were similar to the present study (Tables 2 and 3).

The average Ca²⁺ content was significantly lower (p<0.05) in non-inoculated *Gliricidia* amended SMSC, non-inoculated *Tithonia* amended SMSC and inoculated SMSC without amendment compared to other treatments. A higher Ca²⁺ concentration may obtain due to the availability of CaCO₃ in the SMS.

Figure 4 depicts the Cu²⁺, Fe²⁺ and Zn²⁺ composition in the 100% CAE.




Figure 4: Cu²⁺, Fe²⁺ and Zn²⁺ composition in the 100% CAE (Vertical bars represent standard error (mean ± SE) and mean values designated by the same letter are not significantly different at p<0.05 as determined by DMRT)

Metals significantly inhibit germination and plant growth (Wollan *et al.* 1978). Further, the heavy metal concentration in compost is considered a restrictive factor which is essential to analyze for its safe application as an organic amendment to the soil (Alvarenga *et al.* 2015). Plant micronutrients such as zinc (Zn^{2+}), copper (Cu^{2+}), and iron (Fe^{3+}) had significant differences in treatments in this study. Cu^{2+} was detected only in *Gliricidia* amended SMSC. It was significantly different ($p < 0.05$) between inoculated and non-inoculated while inoculated was higher. The amount of Fe^{3+} was significantly different in most of the treatments. It was significantly higher ($p < 0.05$) in inoculated *Tithonia* amended SMSC followed by its non-inoculated treatment than other treatments. Fe^{3+} content was significantly lower ($p < 0.05$) in inoculated SMSC without amendments followed by non-inoculated urea amended

SMSC. Water extractable Cu^{2+} concentrations in the cattle manure valuing 0.04 mg/L inhibited the root growth of the cress.

In most of the treatments, Zn^{2+} content was significantly different compared to each other. Zn^{2+} content was significantly higher ($p < 0.05$) in inoculated *Gliricidia* amended SMSC followed by inoculated SMSC without amendments. Significantly lower ($p < 0.05$) Zn^{2+} content was recorded at non-inoculated SMSC without amendments. According to Davies (1977) Zn^{2+} contents of SMSC have not exceeded the critical range of 75 to 600 mg/L.

Figure 5 shows the Cr^{2+} and Ni^{2+} composition in the 100% CAE. Potentially toxic heavy metals including chromium (Cr^{2+}), and nickel (Ni^{2+}) were observed in most of the treatments of this study.




Figure 5: Cr^{2+} and Ni^{2+} composition in the 100% CAE (Vertical bars represent standard error (mean \pm SE) and mean values designated by the same letter are not significantly different at $p < 0.05$ as determined by DMRT)

However, Cr^{2+} content was comparatively low. Among the treatments Cr^{2+} content was significantly higher ($p < 0.05$) in non-inoculated *Gliricidia* amended SMSC and lower ($p < 0.05$) in *Tithonia* amended SMSC. Ni^{2+} content was significantly higher ($p < 0.05$) in inoculated *Tithonia* amended SMSC compared to other treatments except inoculated SMSC without amendment. Ni^{2+}

was not detected in *Pueraria* amended SMSC and inoculated urea amended SMSC. Kebrom *et al.* (2019) suggest that the phytotoxic effect in their respective study of them could be due to Ni^{2+} .

Figure 6 depicts the SO_4^{2-} and NH_4^+-N composition in the 100% CAE. NH_4^+-N are possible compounds that generate

phytotoxicity in seed germination (Ramírez *et al.*, 2008).




Figure 6: SO₄²⁻ and NH₄⁺-N composition in the 100% CAE (Vertical bars represent standard error (mean ± SE) and mean values designated by the same letter are not significantly different at p<0.05 as determined by DMRT)

Any of the inorganic forms of nitrogen (ammonia, nitrite and nitrate) in mature compost is of vast importance to determine its quality (Cesaro *et al.* 2019). In the present study, NH₄⁺ content was significantly higher (p<0.05) in inoculated urea amended SMSC and significantly lower (p<0.05) in non-inoculated *Tithonia* amended SMSC and inoculated SMSC without amendments. The highest value of NH₄⁺ concentration in urea may be due to the higher mineralization of amended urea which contains a higher amount of total N% (46%). According to Hoekstra *et al.* (2002), NH₄⁺ of 633 mg/L has been demarcated as a phytotoxic level for compost extracts. However, none of the treatments was found to have this critical level in the present study.

There were no significant differences among most of the treatments in SO₄²⁻ content. However, it was significantly higher (p<0.05) in inoculated urea amended SMSC while significantly lower (p<0.05) in inoculated *Pueraria* amended SMSC. The compost extracts showed a higher amount of SO₄²⁻ may be due to added Epsom salt in the mushroom production substrate.

Pearson correlation coefficients among various parameters of CAEs

A correlation analysis was conducted between GI, RSG, RRG and other chemical parameters of inoculated and non-inoculated CAEs separately, to observe if any relationships between biological and chemical parameters (Tables 2 and 3). They probably indicate the indirect effects of phytotoxicity.

GI showed a higher negative significant correlation (r=-0.841, p<0.01; r=-0.778, p<0.01) with EC in both inoculated and non-inoculated compost samples, respectively. Pearson correlation of pH and Cu was found to be moderately positive and statistically significant (r=0.645, p<0.01) in inoculated treatments while pH and Zn showed a moderately positive and statistically significant (r=0.577, p<0.05) relationship in non-inoculated treatments. In the case of EC, there were positive significant correlations with Cr (r=0.612, p<0.05), Fe (r=0.822, p<0.01), Zn (r=0.828, p<0.01) in inoculated CAEs and with Cr (r=0.625, p<0.05), Fe (r=0.861, p<0.01), Zn (r=0.785, p<0.01), Ni (r=0.517, p<0.05) in non-inoculated CAEs (Table 3). When considering the correlation of

EC and other cations (Na^+ , Ca^{2+} , NH_4^+) and anions (SO_4^{2-}) a positive and significant correlation was observed in both inoculated and non-inoculated aqueous extracts.

On the other hand, in inoculated CAEs, Cr, Fe and Zn showed highly negative and statistically significant relationships ($r=-0.555$, $p<0.05$; $r=-0.634$, $p<0.05$; $r=-0.713$, $p<0.01$) with GI, respectively. Similar relationships were observed in Fe and Zn ($r=-0.766$, $p<0.01$; $r=-0.544$, $p<0.05$) in non-inoculated extracts.

RSG in inoculated CAEs showed a significant negative correlation with SO_4^{2-} ($r=-0.616$, $p<0.05$) while RSG in non-inoculated extracts did not show a significant relationship with any of the parameters.

RRG in inoculated CAEs showed significant negative correlations with EC ($r=-0.887$, $p<0.01$), Cr ($r=-0.521$, $p<0.05$), Fe ($r=-0.717$, $p<0.01$), Zn ($r=-0.738$, $p<0.01$), Na ($r=-0.924$, $p<0.01$), Ca ($r=-0.777$, $p<0.01$), SO_4^{2-} ($r=-0.592$, $p<0.05$), and NH_4^+ ($r=-0.712$, $p<0.01$). RRG in non-inoculated CAEs showed significant negative correlations with EC ($r=-0.725$, $p<0.01$), Fe ($r=-0.734$, $p<0.01$), Na ($r=-0.789$, $p<0.01$), Ca ($r=-0.753$, $p<0.01$), SO_4^{2-}

($r=-0.780$, $p<0.01$), and NH_4^+ ($r=-0.755$, $p<0.01$).

Further, GI in inoculated CAEs showed significant negative correlations with Na ($r=-0.889$, $p<0.01$), Ca ($r=-0.734$, $p<0.01$), SO_4^{2-} ($r=-0.671$, $p<0.01$), and NH_4^+ ($r=-0.717$, $p<0.01$). GI in non-inoculated CAEs showed significant negative correlations with Na ($r=-0.808$, $p<0.01$), Ca ($r=-0.722$, $p<0.01$), SO_4^{2-} ($r=-0.800$, $p<0.01$), and NH_4^+ ($r=-0.807$, $p<0.01$).

EC in inoculated CAEs showed significant positive correlations with Na ($r=0.947$, $p<0.01$), Ca ($r=0.738$, $p<0.01$), SO_4^{2-} ($r=0.639$, $p<0.05$), and NH_4^+ ($r=0.779$, $p<0.01$). EC in non-inoculated CAEs showed significant positive correlations with Na ($r=0.910$, $p<0.01$), Ca ($r=0.648$, $p<0.01$), SO_4^{2-} ($r=0.801$, $p<0.01$), and NH_4^+ ($r=0.892$, $p<0.01$).

Hoekstra *et al.* (2002) found high and significant inverse correlation of EC with RRG and GI similar to the present study. In his study, NH_4^+ was significantly and positively correlated with RRG and GI which was not observed in the present study.

Table 2: Pearson correlation coefficients among various parameter of CAEs treated with in-

	pH	EC	Cu^{2+}	Cr^{2+}	Fe^{3+}	Zn^{2+}	Ni^{2+}	Na^+	SO_4^{2-}	Ca^{2+}	NH_4^+
RSG%	ns	ns	ns	ns	ns	ns	ns	ns	-0.616*	ns	ns
RRG%	ns	-0.887**	ns	-0.521*	-0.717**	-0.738**	ns	-0.924**	-0.592*	-0.777**	-0.712**
GI%	ns	-0.841**	ns	-0.555*	-0.634*	-0.713**	ns	-0.889**	-0.671**	-0.734**	-0.717**
pH	-	ns	0.645**	ns	ns	ns	ns	ns	ns	ns	ns
EC		-	ns	0.612*	0.822**	0.828**	ns	0.947**	0.639*	0.738**	0.779**
Cu^{2+}			-	0.641*	ns	0.558*	ns	ns	ns	ns	ns
Cr^{2+}				-	ns	0.556*	ns	0.524*	ns	ns	0.543*
Fe^{3+}					-	ns	ns	0.768**	0.620*	0.761**	0.792**
Zn^{2+}						-	0.525*	0.766**	ns	ns	ns
Ni^{2+}							-	ns	ns	ns	ns
Na^+								-	0.703**	0.878**	0.805**
SO_4^{2-}									-	0.739**	0.911**
Ca^{2+}										-	0.859**
NH_4^+											-

Note: Values with ** indicates that the correlation is significant at $p<0.01$ level and * indicates that the correlation is significant at $p<0.05$; Blue colour represent the positive correlation and orange colour represent the negative correlation.

Table 3: Pearson correlation coefficients among various parameter of CAEs treated without inoculum

	pH	EC	Cu ²⁺	Cr ²⁺	Fe ³⁺	Zn ²⁺	Ni ²⁺	Na ⁺	SO ₄ ²⁻	Ca ²⁺	NH ₄ ⁺
RSG%	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns
RRG%	ns	-0.725**	ns	ns	-0.734**	ns	ns	-0.789**	-0.780**	-0.753**	-0.755**
GI%	ns	-0.778**	ns	ns	-0.766**	-0.544*	ns	-0.808**	-0.800**	-0.722**	-0.807**
pH	-	ns	ns	ns	ns	0.577*	ns	ns	ns	ns	ns
EC	-	-	ns	0.625*	0.861**	0.785**	0.517*	0.910**	0.801**	0.648**	0.892**
Cu²⁺			-	0.572*	ns	0.581*	ns	ns	ns	ns	ns
Cr²⁺				-	ns	ns	ns	0.637*	0.608*	0.648**	0.744**
Fe³⁺					-	0.598*	0.587*	0.896**	0.834**	0.644**	0.757**
Zn²⁺						-	ns	0.551*	ns	ns	0.630*
Ni²⁺							-	0.599*	ns	ns	ns
Na⁺								-	0.950**	0.874**	0.948**
SO₄²⁻									-	0.914**	0.912**
Ca²⁺										-	0.888**
NH₄⁺											-

Note: Values with ** indicates that the correlation is significant at $p < 0.01$ level and * indicates that the correlations significant at $p < 0.05$; Blue colour represent the positive correlation and orange colour represent the negative correlation.

CONCLUSIONS

Determination of the phytotoxicity in the spent mushroom substrate compost is important for guaranteeing the quality. The seed germination of *R. sativus* in CAEs of different compost mixtures amended with nitrogenous additives and inoculum was not inhibited by any concentrations except non-inoculated *Pueraria* amended SMSC. The higher GI in treatments compared to the control indicated the phyto-stimulant effect. The seed germination of *Tithonia* amended SMSC was not inhibited by any of the concentrations used in this study indicating the higher usability as an amendment in SMS composting. Further, the inoculated SMSC showed higher performance with higher GI compared to non-inoculated SMSC which may be due to the reduction of phytotoxic substances in the compost products. It can be concluded that the SMS integrated with nitrogenous additives and inoculum has the possibility to be deployed as free phytotoxic materials. Large-scale compost generation by SMS and field tests should be conducted as future research to observe crop growth and yield performance.

AUTHOR CONTRIBUTION

SRA conceptualized and designed the study. SRA performed the experiments. SRA and

WMCSJ analyzed and interpreted the data. SRA drafted the manuscript. SRA and WMCSJ critically revised the manuscript.

REFERENCES

- Albuquerque J, De La Fuente C and Bernal M 2011 Improvement of soil quality after “alperujo” compost application to two contaminated soils characterized by differing heavy metal solubility. *Journal of Environmental Management*, 92: 733–741.
- Alvarenga P, Mourinha C, Farto M, Santos T, Palma P, Sengo J, Morais MC and Cunha-Queda C 2015 Sewage sludge, compost and other representative organic wastes as agricultural soil amendments: Benefits versus limiting factors. *Waste Management*, 40: 44–52.
- Amarasinghe SR and Gunawardhana LAASM 2020 Effect of vermicompost with inoculated biochar on Carbon and Nitrogen mineralization in Ultisols and Alfisols. 13th International Research Conference, General Sir John Kotelawala Defense University, 40-45.
- Barral TM and Paradelo R 2011 A Review on the Use of Phytotoxicity as a Compost

- Quality Indicator Dynamic Soil, Dynamic Plant, 2:36-44
- Beyer M (1996). The impact of the mushroom industry on the environment. *Mushroom News* 44(11): 6-13.
- Bolan NS, Kunhikrishnan A and Naidu R 2013 Carbon storage in a heavy clay soil landfill site after biosolid application. *Science of the Total Environment*, 465:216–225.
- Cayuela ML, Millner PD, Meyer SLF and Roig A 2008 Potential of olive mill waste and compost as biobased pesticides against weeds, fungi, and nematodes. *Science of the Total Environment*, 399:11-18.
- Cesaro A, Belgiorno V and Guida M 2015 Compost from organic solid waste: Quality assessment and European regulations for its sustainable use. *Resources, Conservation and Recycling*, 94:72-79.
- Cesaro A, Conte A, Belgiorno V, Siciliano A and Guida M 2019 The evolution of compost stability and maturity during the full-scale treatment of the organic fraction of municipal solid waste. *Journal of Environmental Management*, 232:264-270.
- CHEMetrics 2018 Iron (total, soluble, ferrous)- Phenanthroline Method. 10:1–2.
- Chen M, Xu P, Zeng G, Yang C, Huang D and Zhang J 2015 Bioremediation of soils contaminated with polycyclic aromatic hydrocarbons, petroleum, pesticides, chlorophenols and heavy metals by composting: applications, microbes and future research needs. *Biotechnology Advances*. 33:745–755.
- Davies GR 1977. Pot experiments testing zinc, copper and nickel salts on the growth and composition of crops. *Inorganic Pollution and Agriculture*, Ministry of Agriculture, Fisheries and Food, HMSO, London, 191–204.
- Davis DD, and Fidanza MA 2011 Fresh Recycled Mushroom Compost Suppresses Artillery Fungi Sporulation: A 4-Year Field Study. *Journal of Environmental Horticulture*, 29(2), 91–95.
- Emino ER and Warman PR 2004 Biological assay for compost quality. *Compost Science and Utilization*, 12:342–348.
- Fuentes A, Llorens M, Saez J, Aguilar MI, Ortuno JF and Meseguer VF 2004 Phytotoxicity and heavy metals speciation of stabilised sewage sludges. *Journal of hazard materials*, 108:161–169.
- Gajdos R 1997 Effects of two composts and seven commercial cultivation media on germination and yield. *Compost Science & Utilization*, 5(1):16-37.
- Gariglio NF, Buyatte MA, Pilatti RA, Rossia, DEG and Acosta MR 2002 Use of a germination bioassay to test compost maturity of willow (*Salix* sp.) Sawdust. *New Zealand Journal of Crop and Horticultural Science*, 30 (2):135-139.
- Guo R, Li G, Jiang T, Schuchardt F, Chen T, Zhao Y and Shen Y 2012 Effect of aeration rate, C/N ratio and moisture content on the stability and maturity of compost. *Bioresources Technology*, 112:171–178.
- Hase T and Kawamura K 2012 Evaluating compost maturity with a newly proposed index based on a germination test using *Komatsuna* (*Brassica rapa* var. *peruviridis*) seeds. *J. Mater. Cycles. Waste Management*, 14:220–227.
- Hoekstra NJ, Bosker T, Lantinga EA 2002 Effects of cattle dung from farms with different feeding strategies on germination and initial root growth of cress (*Lepidium sativum* L.). *Agriculture, Ecosystems and Environment* 93, 189-196.
- Huang D, Qin X, Xu P, Zeng G, Peng Z, Wang R, Wan J, Gong X and Xue W 2016 Composting of 4-nonylphenol-contaminated river sediment with inocula of *Phanerochaete chrysosporium*. *BioresourcesTechnology*, 221:47-54
- Iannotti DA, Grebus ME, Toth BL, Madden LV and Hoytink HAJ 1994 Oxygen respirometry to assess the stability and maturity of composted municipal solid

- waste. *Journal of Environmental quality*, 23:1177-1183
- Jain S and Arnepalli DN 2018 Biomineralization as a remediation technique: A critical review. *Geotechnical Characterization and Geo-environmental Engineering Part of the Lecture Notes in Civil Engineering book series (LNCE) 16*: 155–162 DOI: 10.1007/978-981-13-0899-4_19
- Kebrom TH, Woldesenbet S, Bayabil HK, GarciaM, GaoM, Ampim P, Awal R and Fares A 2019 Evaluation of phytotoxicity of three organic amendments to collard greens using the seed germination bioassay. *Environmental Science and Pollution Research*, 26(6):5454–5462.
- Kim D, Lee G, Chang M, Park J, Chung Y, Lee S and Lee T 2011 Purification and Biochemical Characterization of Insoluble Acid Invertase (INAC-INV) from Pea Seedlings. *Journal of Agricultural and Food Chemistry*, 59: 11228–11233. dx.doi.org/10.1021/jf201057c
- Luo Y, Liang J, Zeng G, Chen M, Mo D, Li G, Zhang D 2018 Seed germination test for toxicity evaluation of compost: Its roles, problems and prospects. *Waste Management*, 71:109-114
- Marambe B and Ando T 1992 Phenolic acids as potential seed germination inhibitors in animal waste compost. *Soil Science & Plant Nutrition* 38:727–733.
- Mazumder P, Khwairakpam M and Kalamdhad AS 2020 Bio-inherent attributes of water hyacinth procured from contaminated water body–effect of its compost on seed germination and radicle growth. *Journal of Environmental Management*, 257:109-119.
- Ojuederie OB and Babalola OO 2017 Microbial and Plant-Assisted Bioremediation of Heavy Metal Polluted Environments: A Review. *International Journal of Environmental Research, Public Health*, 14:1504.
- Pampuro N, Santoro E and Cavallo E 2010 Evaluation of maturity and fertilizer capacity of compost derived from swine solid fraction *International Conference on Agricultural Engineering: Towards Environmental Technologies*, 28.
- Peters S, Koschinsky S, Schwieger F and Tebbe CC 2000 Succession of Microbial Communities during Hot Composting as Detected by PCR–Single-Strand-Conformation Polymorphism-Based Genetic Profiles of Small-Subunit rRNA Genes. *Applied and Environmental Microbiology*, 66(3):930–936.
- Ramírez WA, Domene X, Andrés P and Alcañiz JM 2008 Phytotoxic effects of sewage sludge extracts on the germination of three plant species. *Ecotoxicology*, 17:834-844.
- Rynk R, van de Kamp M, Willson GB, Singley ME, Richard TL, Kolega JL, Gouin FR, Laliberty L, Jr Day K, Murphy DW, Hoitink HAJ and Brinton WF 1992 *On-farm composting handbook*, Cornell University Ithaca, New York, USA.
- Said-pullicino D, Erriquens F and Gigliotti GA 2007 Changes in the chemical characteristics of water-extractable organic matter during composting and their influence on compost stability and maturity. *Bioresources Technology*, 98:1822– 1831.
- Siles-castellano AB, Lopez JM, Lopez-Gonzalez JA, Suarez-Estrella F, Jurado MM, Estrella-Gonzales, MJ and Moreno J 2020 Comparative analysis of phytotoxicity and compost quality in industrial composting facilities processing different organic wastes. *Journal of cleaner production*, 252.
- Singh J and Kalamdhad AS 2013a. Assessment of bioavailability and leachability of heavy metals during rotary drum composting of green waste (water hyacinth). *Ecological Engineering*, 52:59–69.
- Singh J and Kalamdhad AS 2013b. Reduction of bioavailability and leachability of heavy metals during vermicomposting

- of water hyacinth. *Environmental Science and Pollution Research*, 20: 8974–8985.
- Sobarzo-Bernal O, Gómez-Merino FC, Alcántar-González G, Saucedo-Veloz C and Libia Trejo-Téllez LI 2021 Biostimulant effects of cerium on seed germination and initial growth of tomato seedlings. *Agronomy*, 11:08.
- SPSS 2006 PASW Statistics for Windows SPSS Inc.
- Stoffella PJ and Kahn BA 2001 Compost utilization in horticultural cropping systems. Boca Raton, FL: Lewis Publishers. 432.
- Tang L, Zeng G, Shen G, Li Y, Zhang Y and Huang D 2008 Rapid detection of picloram in agricultural field samples using a disposable immunomembrane-based electrochemical sensor. *Environmental Science Technology*, 42:1207–1212.
- Wang G, Kong Y, Liu Y, Li D, Zhang X, Yuan J, and Li G 2020 Evolution of phytotoxicity during the active phase of co-composting of chicken manure, tobacco powder and mushroom substrate. *Waste Management*, 114, 25–32.
- Warman PR 1999 Evaluation of seed germination and growth tests for assessing compost maturity. *Compost Science and Utilization*, 7:33-37.
- Wollan E, Davis RD, Jenner S 1978 Effects of sewage sludge on seed germination. *Environmental Pollution*, 17:195-205.
- Wong MH, Cheung YH, Cheung CL 1983 The effects of ammonia and ethylene oxide in animal manure and sewage sludge on the seed germination and root elongation of *Brassica parachinensis*. *Environmental Pollution*, 30:109- 123.
- Yang F, Li GX, Yang QY, Luo WH 2013 Effect of bulking agents on maturity and gaseous emissions during kitchen waste composting. *Chemosphere*, 93:1393–1399
- Zucconi F, Pera A, Forte M and Bertoldi M 1981 Evaluating toxicity of immature compost. *BioCycle*, 2(2):54-57.