





2.

2.1.

2.1.1. Differentiate between primary, secondary, and tertiary sources of information. (15 marks)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

2.1.2. State advantages and disadvantages of tertiary information sources. (15 marks)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

2.1.3. Briefly describe about "patient care Era". (20 marks)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

2.2.

2.2.1. List **five** factors affecting drug dosage.

(10 marks)

.....  
.....  
.....  
.....  
.....

2.2.2. Briefly describe the effect of **one** factor mentioned above 2.2.1 on drug dosage. (15 marks)

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....

2.2.3. Calculate the dose for a child 5 years of age, 39 inches in height and weighing 32 lb, for a drug with an adult dose of 100 mg. (10 marks)

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....

2.2.4. If the recommended dose of vancomycin for a premature infant is 10 mg/kg/day, divided into two equal doses administered every 12 hours, how many milligrams of the drug should be given every 12 hours to an 8.8 lb infant? (15 marks)

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....







