

The momentum distribution of one-dimensional strongly repulsive Bose gas at large momenta

Nandani E.J.K.P.^{1,2,3*} and Xi-Wen Guan.¹

¹*Wuhan Institute of Physics and Mathematics, Chinese Academy of Sciences, Wuhan 430071, People's Republic of China.*

²*University of Chinese Academy of Sciences, Beijing 100049, People's Republic of China.*

³*Department of Mathematics, University of Ruhuna, Matara 81000, Sri Lanka.*

The physics of ultracold Bose gases is a rapidly developing field due to the recent remarkable achievement in the experiment and intensive theoretical investigations. However, when understanding quantum correlations, the calculation of correlation functions of such many-body systems always imposes a big challenge. In this communication, we analytically calculate the momentum distribution of the one-dimensional Bose gas at high momenta through the Fourier transform of the single-particle density matrix. Using the Bethe ansatz wave function of the one-dimensional strongly interacting bosons at the ground state, we find that asymptotic behavior of the momentum distribution for large values of momenta obeys the universal power law C/p^4 decay, where the Tan's contact C can be given explicitly in terms of the generalized exclusion statistics parameter.

Keywords: Bose gas, Momentum distribution, Tan contact.

Acknowledgements: *This work has been supported by CAS_TWAS President's Fellowship for International PhD student.*

**Corresponding Author: nandani@maths.ruh.ac.lk*